



**Europäisches
Patentamt**

**European
Patent Office**

**Office européen
des brevets**

Bescheinigung

Certificate

Attestation

Die angehefteten Unterlagen stimmen mit der ursprünglich eingereichten Fassung der auf dem nächsten Blatt bezeichneten europäischen Patentanmeldung überein.

The attached documents are exact copies of the European patent application described on the following page, as originally filed.

Les documents fixés à cette attestation sont conformes à la version initialement déposée de la demande de brevet européen spécifiée à la page suivante.

Patentanmeldung Nr. Patent application No. Demande de brevet n°

03002993.8

Der Präsident des Europäischen Patentamts;
Im Auftrag

For the President of the European Patent Office

Le Président de l'Office européen des brevets
p.o.

R C van Dijk

THIS PAGE BLANK (USPTO)



Anmeldung Nr:
Application no.: 03002993.8
Demande no:

Anmeldetag:
Date of filing: 11.02.03
Date de dépôt:

Anmelder/Applicant(s)/Demandeur(s):

Universitätsklinikum Charité an
der Humboldt- Universität zu Berlin
Schumannstrasse 20/21
10117 Berlin
ALLEMAGNE

Bezeichnung der Erfindung/Title of the invention/Titre de l'invention:
(Falls die Bezeichnung der Erfindung nicht angegeben ist, siehe Beschreibung.
If no title is shown please refer to the description.
Si aucun titre n'est indiqué se référer à la description.)

Lipid phosphate phosphatases and uses thereof for treating neuronal diseases

In Anspruch genommene Priorität(en) / Priority(ies) claimed /Priorité(s)
revendiquée(s)
Staat/Tag/Aktenzeichen/State/Date/File no./Pays/Date/Numéro de dépôt:

EP/13.09.02/EP 02020679

Internationale Patentklassifikation/International Patent Classification/
Classification internationale des brevets:

G12N9/00

Am Anmeldetag benannte Vertragsstaaten/Contracting states designated at date of
filing/Etats contractants désignées lors du dépôt:

AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HU IE IT LU MC NL
PT SE SI SK TR LI

THIS PAGE BLANK (USPTO)

5 **Lipid phosphate phosphatases and uses thereof for treating neuronal diseases**

The present invention relates to lipid phosphate phosphatase proteins, genes coding for them, vectors and cells comprising them, antibodies directed against them, methods of identifying compounds binding to them and functional interactors as well as to the use of proteins, genes,
10 vectors, cells, interacting compounds and functional interactors for treating neuronal diseases and/or injuries.

Axons in the central nervous system (CNS) elongate through the extracellular space over long distances (N. Tessier-Lavigne and C. S. Goodman (1996) Science 274:1123-1133). This oc-
15 curs during development (C. S. Goodman (1996) Annu. Rev. Neurosci. 19:341-377 and H. Super and E. Soriano (1994) J. Comp. Neurol. 344:101-120) and during axonal sprouting in response to partial deafferentation (C. Cotman et al. (1977) J. Neurocytol. 6:455-464 and M. Frotscher et al. (1997) Trends Neurosci. 20:218-223). The extracellular space, however, is an
20 outgrowth repellent environment that allows axonal elongation only under specific molecular conditions (E. Stein and N. Tessier-Lavigne (2001) Science 291:1928-1938). Molecules in-
25 volved in axonal outgrowth, such as semaphorins, netrins, or ephrins (S. A. Colamarino and M. Tessier-Lavigne (1995) Cell 81:621-629, H Kobayashi et al. (1997) J. Neurosci. 17:8339-8352, E. Stein et al. (1999) J. Neurosci. 19:8585-8893 and A. Steup et al. (2000) Mol. Cell Neurosci. 15:141-155) are able to transduce outgrowth promoting as well as inhibiting signals
to elongating axons via specific receptors.

In the hippocampus, afferent connections from the entorhinal cortex enter in a layer-specific manner during development (T. Skutella and R. Nitsch (2001) Trends Neurosci. 24:107-113). This specific axonal navigation depends on molecular cues expressed along the pathway and
30 in the target region (T. Skutella and R. Nitsch, *supra*) Transection of entorhinal axons in the adult leads to a specific deafferentation in the hippocampus with subsequent regenerative axon sprouting by remaining afferents into the denervated zones (C. Cotman et al, *supra* and D.A Matthews et al. (1976) Brain Res. 115:23-41). It has been shown that signaling via bio-
35 active lipid phosphates such as phosphatidate (PA), lysophosphatidate (LPA) or sphingosine-1-phosphate (S-1-P) are involved in cell migration, mitogenesis and neurite retraction (K.

Jalink et al. (1994) Biochim. Biophys. Acta 1198:185-196, W. H. Moolenaar (1995) Curr. Opin. Cell Biol. 7:203-210 and N. Zhang et al (1997) Nature 385:64-67) and in particular it has been shown that signaling via extracellular LPA plays an important role in CNS development and that postmitotic neurons are at least one endogenous source for LPA in the nervous system (N. Fukushima et al (2000) Dev. Biol. 228:6-18). LPA has properties of an extracellular neurite repellent factor (K. Jalink (1994) *supra* and K. Jalink et al (1993) Cell Growth Differ. 4:247-255). It is present in the extracellular space of the nervous system (Fukushima et al. (2000) *supra* and J. Bothmer et al (1992) Neurochem. Int. 21:223-228) and mediates diverse cellular responses through the activation of multiple signal transduction pathways (W. H. Moolenaar (1995) *supra*). One major structural effect of LPA on neurons is rapid neurite retraction with subsequent cell rounding. Therefore, LPA and similar bioactive lipid phosphatases inhibit a regrowth of axons following neuronal lesion. Therefore, it is a problem known in the art, that after neuronal damage due to, for example, neuronal disease or trauma a regrowth of axons does not occur.

Within the context of the present invention it has been found that the expression of a family of genes called plasticity-related genes (PRGs) overcomes the repellent effect of bioactive lipid phosphates, in particular of LPA and, thus, allows the regrowth of axons in spite of the presence of bioactive lipid phosphates. Therefore, the present invention is directed at an isolated protein comprising the same or substantially the same amino acid sequence selected from the group consisting of human PRG-1, human PRG-2, mouse PRG-1, mouse PRG-2 and rat PRG-1 (depicted in SEQ ID NOs: 1 to 5), respectively, or a splice variant or a salt thereof. A protein having substantially the same amino acid sequence comprises proteins with at least about 95%, preferably at least about 96%, more preferably at least about 97%, more preferably with at least about 98% and most preferably with at least about 99% amino acid sequence identity. The amino acid exchanges are preferably so called conservative changes meaning substitutions of, for example, a polar amino acid residue by another polar amino acid residue, of an acidic amino acid residue by another acidic amino acid residue or of a basic amino acid residue by another basic amino acid residue.

Proteins having substantially the same amino acid within the meaning of this invention exhibit in a preferred embodiment lipid phosphate phosphatase activity. The lipid phosphate phosphatase activity of a given protein with substantially the same amino acid can be tested, for example, by the ectophosphatase assay described in example 11 below. The proteins em-

ployed in the assay can either be purified from cells or can be recombinantly expressed and purified by methods well known in the art.

5 In one embodiment of the present invention the protein comprises at least one fragment of the human PRG-1 and PRG-2 or mouse PRG-1 and PRG-2 or rat PRG-1. A fragment within the meaning of the present invention refers to one of the proteins according to SEQ ID NOs: 1 to 5 bearing at least one N-terminal, C-terminal and/or internal deletion. The resulting fragment has a length of at least about 50, preferably of at least about 100, more preferably of at least about 150, more preferably of at least about 200, more preferably of at least about 250, more
10 preferably of at least about 300, more preferably of at least about 350 and most preferably of at least about 400 amino acids.

Preferably the fragment is an N-terminal fragment which comprises 330 amino acids or less as outlined above, which are highly conserved between, for example, PRG-1 and members of
15 the family of LPP membrane-associated phosphatic acid phosphatase ecto-enzymes, which have six membrane spanning domains with their active site located on the external surface of the plasma membrane. This domain comprises preferably the catalytic region. For example, human PRG-1 carries a catalytic histidine at position 252, which is involved in the phosphatase activity of human PRG-1. Therefore, in a preferred embodiment any N-terminal fragment
20 of the proteins of the present invention comprises the catalytic site including the conserved His-residue. The fragment itself has preferably an amino acid sequence identity with hPRG-1, hPRG-2, mPRG-1, mPRG-2 and rat PRG-1, respectively, of at least about 95%, preferably of at least about 96%, more preferably of at least about 97%, more preferably of at least about 98%, more preferably of at least about 99% and most preferably of 100%.

25 The C-terminal cytoplasmatic part of the PRG proteins is potentially involved in regulation of lipid phosphate phosphatase activity and/or signaling and, thus, a further preferred fragment comprises a C-terminal fragment, which comprises about 413 amino acids or less as outlined above and which comprises regions required for above activity of the PRG proteins. The
30 fragment itself has preferably an amino acid sequence identity with hPRG-1, hPRG-2, mPRG-1, mPRG-2 and rat PRG-1, respectively, of at least about 95%, preferably of at least about 96%, more preferably of at least about 97%, more preferably of at least about 98%, more preferably of at least about 99% and most preferably of 100%.

In a further aspect the present invention is directed at a nucleic acid, which comprises at least one nucleic acid encoding one of the proteins of the present invention. Preferably the nucleic acid consists of DNA or RNA, wherein the DNA preferentially is either single or double stranded. Also comprised are DNA's, which hybridize to one of the aforementioned DNA's under stringent conditions like, for example, hybridization at 60°C in 2.5 x SSC buffer and several washes at 37°C at a lower buffer concentration like, for example, 0,5 x SSC buffer and which encode proteins exhibiting lipid phosphate phosphatase activity and/or association with plasma membranes. Additional reagents required for carrying out stringent Northern or Southern blots like, for example, single stranded salmon sperm DNA are well known in the art.

In a preferred embodiment of the nucleic acid of the present invention the nucleic acid comprises a nucleic acid selected from the group consisting of the human PRG-1 gene, the human PRG-2 gene, the mouse PRG-1 gene, the mouse PRG-2 gene and the rat PRG-1 gene (see SEQ ID NOs: 6 to 10).

In a further embodiment the nucleic acid of the present invention further comprises at least one promoter, enhancer, intron and/or polyA-sequence. Preferred promoters or enhancers possess tissue specificity, in particular neuronal specificity and more particular a specificity for growing neurons. Examples of such promoters and/or enhancers are the neuron specific enolase promoter (Erickson, R.P. and Bernard, O. (2002) J. Neuro Science Res 68:738-44), the peripherin promoter (Weinstein, D.E. et al. (1999) Brain Res. Dev. Brain Res. 116:29-39), the synapsin promoter (Flood, D.G. et al. (1999) Am. J. Pathol. 155:663-72) and the Thy 1 promoter (Kahle, P. J. et al. (2001) Am. J. Pathol. 159:2215-25).

In some instances it might be desirable to interfere with, for example, the transcription or translation of the nucleic acids of the present invention and, therefore, the present invention is also directed at a nucleic acid, which is complementary to the nucleic acid of the present invention and, thus, is capable of inhibiting, for example, transcription or translation. A preferred embodiment of such a complementary nucleic acid is a so called anti-sense oligonucleotide (R. Q. Zheng and D. M. Kemeny (1995) Clin. Exp. Immunol. 100:380-2, W. Nellen and C. Lichtenstein (1993) Trends. Biochem. Sci. 18:419-423 and C. A. Stein (1992) Leukemia 6:967-74), ribozymes (M. Amarzguoui and H. Prydz (1998) Cell. Mol. Life Sci. 54:1175-1202, N. K. Vaish et al (1998) Nucleic Acids Res. 96:5237-5242, Persidis (1997)

Nat. Biotechnol. 15:921-922 and L. A. Couture and D. T. Stinchcomb (1996) Trends Genet. 12:510-515) and/or so called small interfering RNA-molecules (siRNAs) (S. M. Elbashir et al. (2001) Nature 411:494-498). Anti-sense oligonucleotides are able to decrease the stability of the above described nucleic acids and/or can inhibit the translation. Similarly the use of
5 siRNA-oligonucleotides also leads to a reduction in the amount of the translated polypeptides. The anti-sense oligonucleotides have in a preferred embodiment a length of at least 20, preferable of at least about 30, more preferably of at least about 40 and most preferably a length of at least about 50 nucleic acids.

10 Oligonucleotides are generally rapidly degraded by endo- or exonucleases, which are present in the cell, in particular by DNases und RNases and, therefore, it is advantageous to modify the nucleic acids which are used, for example, in anti-sense strategies, as ribozymes or siRNAs to stabilize them against degradation and thereby prolong the time over which an effective amount of the nucleic acid is maintained within the cell (L. Beigelmann et al. (1995) Nu-
15 cleic acids Res. 23:3989-94, WO 95/11910, WO 98/37340 and WO 97/29116). Typically such a stabilization can be obtained by the introduction of one or more internucleotide phosphate groups and/or by the introduction of one or more non-phosphor-internucleotides.

Suitable modified internucleotides are summarized in, for example, Uhlmann and Peimann
20 (1990) Can. Rev. 90:544. Modified internucleotide phosphate residues and/or non-phosphate bridges which can be used in a nucleic acid of the invention comprise, for example, methylphosphonate, phosphorthioate, phosphoramidate, phosphordithionate, phosphate ester, non-phosphor internucleotide analogues, which can be used in nucleic acids of the invention include, for example, siloxane bridges, carbonate bridges, carboxymethylester, acetamid bridges
25 and/or thio ether bridges.

A further aspect of the present invention is directed at a vector comprising a protein according to the present invention and/or a nucleic acid according to the present invention. A vector within the meaning of the present invention is a protein or a nucleic acid or a mixture thereof
30 which is capable of being introduced or of introducing the proteins and/or nucleic acid comprised into a cell. It is preferred that the proteins encoded by the introduced nucleic acid are expressed within the cell upon introduction of the vector.

In a preferred embodiment the vector of the present invention comprises plasmids, phagemids, phages, cosmids, artificial mammalian chromosomes, knock-out or knock-in constructs, viruses, in particular adenovirus, vaccinia virus, lentivirus (Chang, L.J. and Gay, E.E. (2000) *Curr. Gene Therap.* 1:237-251), Herpes simplex virus (HSV-1, Carlezon, W.A. et al. (2000) *Crit. Rev. Neurobiol.*), baculovirus, retrovirus, adeno-associated-virus (AAV, Carter, P.J. and Samulski, R.J. (2000) *J. Mol. Med.* 6:17-27), rhinovirus, human immune deficiency virus (HIV), filovirus and engineered versions thereof (see, for example, Cobinger G. P. et al (2001) *Nat. Biotechnol.* 19:225-30), virosomes, "naked" DNA liposomes, and nucleic acid coated particles, in particular gold spheres. Particularly preferred are viral vectors like adeno-viral vectors or retroviral vectors (Lindemann et al. (1997) *Mol. Med.* 3:466-76 and Springer et al. (1998) *Mol. Cell.* 2:549-58). Liposomes are usually small unilamellar or multilamellar vesicles made of cationic and anionic lipids, for example, by ultrasound treatment of liposomal suspensions. The DNA can, for example, be ionically bound to the surface of the liposomes or internally enclosed in the liposome. Suitable lipid mixtures are known in the art and
15 comprise, for example, DOTMA (1, 2-Dioleoyloxypropyl-3-trimethylammoniumbromid) and DPOE (Dioleoylphosphatidylethanolamin) which both have been used on a variety of cell lines.

Nucleic acid coated particles are another means for the introduction of nucleic acids into cells
20 using so called "gene guns", which allow the mechanical introduction of particles into the cells. Preferably the particles itself are inert, and therefore, are in a preferred embodiment made out of gold spheres.

In a further aspect the present invention is directed at an isolated cell comprising a protein of
25 the present invention, a nucleic acid of the present invention and/or a vector of the present invention. Cells of the present invention can be prokaryotic or eukaryotic cells and in a preferred embodiment the cells of the present invention are stem cells, in particular non-human embryonic stem cells, embryonic stem cell lines, foetal stem cells, adult stem cells, neuronal precursor cells or neuronal cells in particular axons (Hsich, G. et al. (2002) *Hum. Gene*
30 *Therap.*, 13:579-604 and Martinez-Serrano, A. et al. (2001) *Curr. Gene Therap.* 1:279-299).

A further aspect of the present invention is a transgenic non-human animal generated from a cell of the present invention. The animal can be a mosaic animal, which means that only part of the cells making up the body comprise cells of the present invention or all cells of the ani-

mal can be derived from a cell of the present invention and those animals can be either homo- or heterozygous with respect to the nucleic acid of the present invention contained within the cell of the present invention. In a preferred embodiment the transgenic animals are either homo- or heterozygous knock-out or knock-in animals with respect to the genes which code for the proteins of the present invention.

In a further aspect the present invention is directed at an antibody directed against a protein of the present invention. The term "antibody" comprises monoclonal and polyclonal antibodies and binding fragments thereof, in particular Fc-fragments as well as so called "single-chain-antibodies" (Bird R. E. et al (1988) Science 242:423-6) and diabodies (Holliger P. et al (1993) Proc. Natl. Acad. Sci. U.S.A. 90:6444-8).

In a further aspect the present invention is directed at a method of producing a protein of the present invention or a nucleic acid of the present invention and comprises the steps of: a) cultivating a cell of the present invention and b) isolating the protein and/or the nucleic acid. If the method is used primarily to isolate nucleic acids then in an preferred embodiment the cells, which are used are prokaryotic cells, in particular *E. coli*. cells If the method is used primarily for the isolation of proteins of the invention than the cells can be either of prokaryotic or eukaryotic origin. Someone of skill in the art is aware of a variety of different cell types suitable for the production of proteins like, for example, *E. coli*, Sf9, Hi5, *P. pastoris* and HeLa. Eukaryotic cells are preferably chosen, if it is desired that the proteins produced by the cells exhibit an essentially natural pattern of glycosylation and prokaryotic cells are chosen, if, for example, glycosylation or other modifications, which are normally introduced into proteins only in eukaryotic cells, are not desired or not needed.

In a further aspect the present invention is directed at a method of isolating compounds interacting with a protein of the present invention comprising the steps of: a) contacting one or more of the proteins of the present invention, preferably one, with at least one potentially interacting compound, and b) measuring binding of said compound to said protein. This method is suitable for the determination of compounds that can interact with the proteins of the present invention and to identify, for example, inhibitors, activators, competitors or modulators of proteins of the present invention, in particular inhibitors, activators, competitors or modulators of the enzymatic activity of the proteins of the present invention.

The potentially binding substance, whose binding to the protein of the present invention is to be measured, can be any chemical substance or any mixture thereof. For example, it can be a substance of a peptide library, a combinatorial library, a cell extract, in particular a plant cell extract, a "small molecular drug", a protein and/or a protein fragment.

5

The term "contacting" in the present invention means any interaction between the potentially binding substance(s) with the proteins of the invention, whereby any of the two components can be independently of each other in a liquid phase, for example in solution, or in suspension or can be bound to a solid phase, for example, in the form of an essentially planar surface or
10 in the form of particles, pearls or the like. In a preferred embodiment a multitude of different potentially binding substances are immobilized on a solid surface like, for example, on a compound library chip and the protein of the present invention is subsequently contacted with such a chip.

15 The proteins of the present invention employed in a method of the present invention can be full length proteins or a fragments with N/C-terminal and/or internal deletions. Preferably the fragments are either N-terminal fragments comprising the enzymatic region of the protein or C-terminal fragments comprising the cytoplasmic region, depending on whether potentially interacting compounds are sought that specifically interact with the N- or C-terminal fragment

20

Measuring of binding of the compound to the protein can be carried out either by measuring a marker that can be attached either to the protein or to the potentially interacting compound. Suitable markers are known to someone of skill in the art and comprise, for example, fluorescence or radioactive markers. The binding of the two components can, however, also be
25 measured by the change of an electrochemical parameter of the binding compound or of the protein, e.g. a change of the redox properties of either the protein or the binding compound, upon binding. Suitable methods of detecting such changes comprise, for example, potentiometric methods. Further methods for detecting and/or measuring the binding of the two components to each other are known in the art and can without limitation also be used to
30 measure the binding of the potential interacting compound to the protein or protein fragments of the present invention. The effect of the binding of the compound or the activity of the protein can also be measured indirectly, for example, by assaying the phosphatase activity of the protein after binding.

As a further step after measuring the binding of a potentially interacting compound and after having measured at least two different potentially interacting compounds at least one compound can be selected, for example, on grounds of the measured binding activity or on grounds of the detected increase or decrease of protein activity, in particular lipid phosphate phosphatase activity upon binding. The phosphatase activity can be measured, for example, as described in example 11.

The thus selected binding compound is then in a preferred embodiment modified in a further step. Modification can be effected by a variety of methods known in the art, which include without limitation the introduction of novel side chains or the exchange of functional groups like, for example, introduction of halogens, in particular F, Cl or Br, the introduction of lower alkyl groups, preferably having one to five carbon atoms like, for example, methyl, ethyl, n-propyl, isopropyl, n-butyl, isobutyl, tert-butyl, n-pentyl or iso-pentyl groups, lower alkenyl groups, preferably having two to five carbon atoms, lower alkynyl groups, preferably having two to five carbon atoms or through the introduction of, for example, a group selected from the group consisting of NH₂, NO₂, OH, SH, NH, CN, aryl, heteroaryl, COH or COOH group.

The thus modified binding substances are then individually tested with the method of the present invention, i.e. they are contacted with the protein and subsequently binding of the modified compounds to the protein is measured. In this step both the binding per se can be measured and/or the effect of the function of the protein like, e.g. the enzymatic activity of the protein can be measured. If needed the steps of selecting the binding compound, modifying the binding compound, contacting the binding compound with a protein of the invention and measuring the binding of the modified compounds to the protein can be repeated a third or any given number of times as required. The above described method is also termed "directed evolution" since it involves a multitude of steps including modification and selection, whereby binding compounds are selected in an "evolutionary" process optimizing its capabilities with respect to a particular property, e.g. its binding activity, its ability to activate, inhibit or modulate the activity, in particular the phosphatase activity of the proteins of the present invention.

A further aspect of the present invention is a method of isolating compounds functionally interacting with the activity of the proteins of the present invention comprising the steps of: a) contacting a neuronal cell that comprises a wt nucleic acid coding for a protein selected from

the group consisting of SEQ ID NOs: 1 to 5, a splice variant thereof, or a fragment thereof with a potential functional interactor, b) contacting the cell with a bioactive lipid phosphate, and c) measuring neurite movement or phosphatase activity.

5 The term “contacting” has to be understood as previously defined and comprises any possibility of interaction between the potential functional interactor and a neuronal cell. Contacting also comprises the introduction of the potential functional interactor into the neuronal cell which can be effected by a variety of methods including, for example, electroporation, which allows influx of a potential functional interactor contained in the medium surrounding the
10 neuronal cell into the neuronal cell. A neuronal cell that comprises a wt nucleic acid coding for a PRG protein (as indicated in SEQ ID NOs: 1 to 5) can be any neuronal cell capable of neurite movement. The cell may or may not express the wild-type nucleic acid depending on, for example, the developmental stage of the neuronal cell. For example entorhinal cortex cells of embryonic day 16 (E16), which exhibit neurite movement, do not express PRG-1 while
15 entorhinal cortex cells of postnatal day 0 (P0) do express PRG-1. The choice of either a cell that already expresses or does not express the wild-type nucleic acids will depend on the functional interaction of the potential functional interactor that is sought. If, for example, a functional interactor is sought, that activates transcription within the cell normally not expressing PRG-1 than, for example, an E16 cell could be chosen. If functional interactors are
20 sought that primarily interact on the protein level, i.e. that activate or suppress phosphatase activity of already expressed PRG-1 than neuronal cells expressing PRG-1 would be chosen like, for example, P0 cells.

Bioactive lipid phosphates are lipid phosphates which inhibit neurite movement of neuronal
25 cells, which do not express PRG-1 cells, like for example, E16 cells. Examples of such bioactive lipid phosphates comprise PA, LPA and S-1-P. Whether a lipid phosphate, which can be used in the method of the present invention is bioactive can be determined by, for example, the experiment described in Example 9.

30 Methods for measuring neurite movements are well-known in the art and are described, for example, in Savaskan et al. (1999) European J. Neurosci. 11:319-326. One way of measuring neurite movement is scoring of the cells after contacting with a bioactive lipid phosphate. It is possible to categorize the cells into at least three different categories, i.e. round cells, cells with short processes and cells with long processes. It is also possible to quantify the effect by

measuring the length of the neurite processes. The effect of the potential functional interactor can be determined by comparing the neurite movement of the neuronal cell, the categories the cells are in or the length of the neurite processes after contacting the cell with a bioactive lipid phosphate with or without the functional interactor. Other ways to assess the effect of potential functional interactors are the determination of the expression level of the PRG genes and/or proteins or the enzymatic activity of the proteins.

In a further embodiment the method includes the additional steps of: a) contacting a neuronal cell that comprises a mutant nucleic acid coding for a mutant of the proteins selected from the group consisting of SEQ ID NOs: 1 to 5, a splice variant thereof, or a fragment thereof or that contains a knock-out of the wt nucleic acid coding for one of said proteins with a potential functional interactor, b) contacting said cell with a bioactive lipid phosphate, and c) measuring neurite movement. The above described neuronal cell is preferably incapable of expressing a functional PRG protein and, thus, can not be stimulated by any functional interactor to activate PRG genes or PRG protein function. Therefore, any potential functional interactor, which shows an effect on the neurite movement of neuronal cells comprising wild-type PRG genes but shows no effect in the neuronal cell comprising mutants or knock-out PGR genes have thereby been shown to functional interact with the PRG genes. Once such an interactor has been identified the mode of functional interaction can be further analyzed and to that end the amount of PRG mRNA and/or protein expressed or the activity of the PRG protein can be determined by a variety of different techniques, which are known in the art.

In a preferred embodiment the method of the invention comprises the further steps of: a) modifying the functional interactor to generate a variety of modified functional interactors, b) contacting a neuronal cell comprising a wild-type nucleic acid coding for a protein selected from the group consisting of SEQ ID NOs: 1 to 5, a splice variant thereof, or a fragment thereof and if needed a cell that comprises a mutant nucleic acid coding for a mutant of the protein selected from the group consisting of SEQ ID NOs: 1 to 5, a splice variant thereof, or a fragment thereof with the modified functional interactors, c) contacting said cell or cells with a bioactive lipid phosphate, d) measuring the neurite movement, and e) if needed repeating steps a) to d) for one or more times. The modification of the functional interactor can be any of the modifications outlined above with respect to the modification of an interacting compound and the modification and selection steps can be repeated one or several times until

a functional interactor has been selected that shows the desired functional interaction, e.g. repression or activation of the activity in particular of the enzymatic activity of PRG proteins.

5 In a further embodiment of the method of the present invention the interacting compound identified as outlined above or the functional interactor identified as outlined above, which may or may not have gone through additional rounds of modification and selection, is admixed with suitable auxiliary substances and/or additives. Such substances comprise pharmacological acceptable substances, which increase the stability, solubility, biocompatibility, or biological half-life of the interacting compound or the functional interactor or comprise substances or materials, which have to be included for certain routes of application like, for example, intravenous solution, sprays, Band-Aids or pills.

10 According to a further aspect of the present invention a pharmaceutical composition is provided for the treatment of neuronal injuries or diseases comprising a protein of the invention, a nucleic acid of the invention, a vector of the invention, a cell of the invention, an antibody of the invention, a binding component isolated by a method of the invention and/or a functional interactor isolated by a method of the invention and if needed suitable auxiliary substances and/or additives.

20 Accordingly, a further aspect of the present invention is the use of a pharmaceutical composition of the invention for the production of a medicament for the treatment of neuronal diseases or injuries. Neuronal diseases which can be treated with the pharmaceutical composition comprise spinal cord lesion, Alzheimer disease and stroke. Typical neuronal injuries comprise traumata of any sort in particular head traumata resulting in the damage of neurons and in particular the severing of neuronal connections.

25 As stated above it has also been found that PRG proteins are differentially expressed in certain tissues and that they have been found to be differentially expressed in certain diseases, however, differential expression is also associated with certain disease states. Thus, PRG proteins present attractive targets for diagnosis and treatment of a variety of diseases. Therefore, another aspect of the present invention is the use of the proteins or nucleic acids of the present invention as diagnostic marker for the diagnosis of a disease or disease state, whereby the presence, the absence, or the amount of PRG proteins is evaluated by, for example, im-

munological methods, RT-PCR, Northern blot. For the immunological detection and/or quantification methods the antibodies of the present invention can be used.

5 As PRG is differentially regulated in neuronal diseases PRG proteins or nucleic acids are in a preferred embodiment used as diagnostic marker for the diagnosis of neuronal diseases.

10 Furthermore PRG proteins are overexpressed in tumors. PRG-1, for example, is overexpressed in a variety of tumor cells, e.g. astroglioma WHO Grad III-IV, neuroblastoma, kidney cell carcinoma, myoblastoma and ovarian cell carcinoma, and in particular it has also been found to be overexpressed in migrating, i.e. metastasizing cancer cells which have lost the anchorage dependence for growth. Therefore, PRG proteins also present attractive targets for diagnoses and treatment of tumors, in particular metastasizing tumors. Consequently, a further aspect of the present invention is the use of PRG proteins and nucleic acids of the present invention as a tumor markers and, preferably, as metastatic markers. Because of the above property PRG proteins are also therapeutic targets for the development of drugs, which modulate, preferably inhibit the function of PRG proteins. Such drugs can be identified with above-described methods for identifying interacting or functional interacting compounds.

20 In a preferred embodiment, PRG proteins of the present invention are used for diagnosis of cancers selected from the group of cancers consisting of neuroblastoma, astroglioma, ovarian cell carcinoma, prostatic cell carcinoma and breast cell carcinoma.

25 In addition, PRG proteins, in particular PRG-1, have been found to be overexpressed in differentiating sperm cells and, thus, a further embodiment of the present invention is the use of the PRG proteins or nucleic acids as diagnostic targets for the diagnosis of infertility, in which a lack of PRG proteins would indicate either a low amount of differentiating sperm cells or the malfunction of the sperm cells. The amount of expression of PRG proteins or nucleic acids can be detected by, for example, *in situ* immunofluorescence, *in situ* Northern blots. The skilled persons knows a variety of additional methods that are suitable to determine the amount and distribution of PRG proteins and RNA within a cell and/or tissue.

30 The following figures and examples merely serve to illustrate the invention and should not be construed to restrict the scope of the invention to the particular embodiments of the invention

described in the examples. All references cited in the text and the disclosure of the priority application EP 02 020 679.3 are hereby incorporated in their entirety by reference.

Figures

5

Fig. 1 Human PRG-1 amino acid sequence (SEQ ID NO. 1). The first 300 amino acids are highly conserved among LPP family members. The other 400 amino acids (gray boxed sequence) of PRG-1 show no homologies to known sequences. The catalytic histidine (His-252) conserved in all members of the LPP-super family is marked with an asterisk.

10

Fig. 2 Hydrophobicity profile of PRG-1 protein predicted by the Kyte and Doolittle algorithms. The numbers at the bottom of the profile refer to amino acid residues from the amino terminus. The gray shaded area of PRG-1 is predicted as hydrophilic and located in the cytosol.

15

Fig. 3 Northern blot analysis of PRG-1 mRNA expression

Fig. 4 Expression pattern of PRG-1 mRNA in the developing and lesioned rat brain detected by *in situ* Northern blot. Panel A shows the Toluidene blue staining of a brain section on embryonic day 16 (E16). Panel B shows the *in situ* hybridization signal with a probe specific for PRG-1 mRNA of the same section as shown in panel A. Panel C shows a Toluidene blue staining of a brain section on embryonic day 19 and panel D shows the *in situ* hybridization with a probe specific for PRG-1 mRNA of the same section as shown in panel C. Panels E through K show *in situ* Northern blot analysis of brain sections at days 0, 5, 10, 15, 30 after birth, in an adult and one day after lesion, respectively. The scale bar in E19 equals 850 μ m. The scale bar in P30 equals 740 μ m and also applies to P0-P15. The scale bar in 1dal equals 500 μ m and also applies to adult. "LV" means lateral ventricle, "LP" means lateral posterior thalamic nucleus, "LD" means laterodorsal thalamic nucleus, "bcp" means basal telencephalic plate, posterior part, "CA1" means cornu ammonis, "DG" means dentate gyrus, "RSG" means retrosplenial granular cortex, "dal" means days after lesion.

20

25

30

- Fig. 5 Quantification of the *in situ* hybridization signals in different regions of the brain after lesion. "gcl" means granule cell layer, "CA3" means cornu ammonis 3, "CA1" means cornu ammonis and "dal" means days after lesion.
- 5 Fig. 6 Overexpression of a PRG-1-eGFP fusion protein in COS-7 cells. Panel A shows the fluorescence of the green fluorescent protein, panel B shows the fluorescence of a fluorescently labeled anti-PRG-1 peptide antibody and panel C shows the co-localization of the fluorescence of the PRG-1-eGFP and the anti-PRG-1 antibody. In each picture the processes of the COS-7 cells are marked with a white arrow. The scale bar depicts a length of 10 μ m. Panel D shows the result of a Western blot using an antiserum raised against a C-terminal peptide of PRG-1.
- 10
- Fig. 7 Immunocytochemical analysis of PRG-1 in the adult rat hippocampus prior and after lesion. Pyramidal neurons are labeled in the CA1 and CA3 region. Polymorphic cells are stained in the hilus. The outer molecular layer is marked with black errors. "Gcl" means granule cell layer "Oml" means outer molecular layer and "hi" means hilus. The scale bar in panel A and B equals 580 μ m. Panel C shows immunoblots from total protein extracts of adult control and deafferented hippocampus five days after lesion.
- 15
- 20
- Fig. 8 Panel A shows a higher magnification of the boxed area of panel B of Fig. 7. Immuno-stained axons are marked with black arrows and the terminal branches with white arrows. Panel B shows an electron micrograph of a PRG-1 immunopositive axon. The immunopositive axon is delineated by black arrows, while its terminal branch is delineated by gray arrows. "oml" means outer molecular layer and "ax" means axon and "s" means spine. The scale bar in panel A equals 20 μ m and the scale bar in panel B equals 0.4 μ m.
- 25
- Fig. 9 Panel A-C show the result of an RT-PCR analysis of RNA from tissues from E16 and P0 explants. The amplification was carried out with primers specific for PRG-1 (panel A), EDG 2, 4 or 7 (panel B) and β -actin (panel C). Panel D and E depict explants from rat entorhinal cortex at embryonic day 16 and panel F and G depict rat entorhinal cortex at postnatal day 0 (bottom row of panels). Outgrowing axons are marked with white arrows and the panels D and F show neurite retraction in the
- 30

presence of vehicle (0,9% NaCl) while panels E and G show neurite retraction upon the addition of 100 nmol/l LPA. The scale bar in panel G equals 20 μ m.

Fig. 10 Panel A shows the neurite outgrowth length in control cultures (left bar) and LPA treated cultures, (right bar) in E16 and postnatal explants (P0) in μ m. Panel B shows the dose-response of LPA on P0 and E16 explants.

Fig. 11 Depicts cell rounding and neurite retraction in response to LPA in N1E-115 cells. The cells depicted in the first three top panels were transfected with a plasmid containing only eGFP and were not treated with LPA but with vehicle only. The cells in the second row were transfected with a plasmid containing only eGFP but were treated with 10 μ mol/l LPA, the cells of the third row were transfected with a plasmid coding for a PRG-1-eGFP fusion protein and were treated with 10 μ mol/l LPA while the cells depicted in the bottom row of panels were transfected with a mutant PRG-1-eGFP fusion protein, which carried a His-Lys exchange in the catalytic histidine of PRG-1 (PRG-1^{His/Lys}) and were also treated with 10 μ mol/l LPA. The panels on the left show transfected cells, panels in the middle show nuclear staining (Hoechst staining) and panels on the right show merged images with f-actin staining.

Fig. 12 Panel A shows the quantification of the results shown in Fig. 11 and of a similar experiment performed with wild type N1E-115 cells (wt). Panel B shows the results of RT-PCR using LPA receptor specific primers to assess LPA receptor expression in wild type N1E-115 cells (wt) or PRG-1 transfected N1E cells (PRG-1). Panel C shows the results of RT-PCR using β -actin specific primers to assess β -actin expression in wt N1E-115 cells and in PRG-1 transfected cells. Panel D shows a dose response of LPA treatment on neurite length of PRG-1 overexpression cells. Panel E shows cell viability of N1E-115 cells 48-72 h after transfection. Panel F shows the phosphatase activity of intact cells overexpressing eGFP, PRG-1-eGFP, and PRG-1^{His/Lys}.

Fig. 13 Schematic diagram of the proposed axon growth mechanism in a lipid phosphate lipid rich environment. "R" means receptor mediating a retraction signal.

Fig. 14 Depicts the proposed structure of the human PRG-1 wherein those parts presumed to be arranged on the extracellular surface inserted in the plasma membrane and protruding into the cytoplasm are shown.

5

Examples

1. Isolation of PRG-1

Animals and surgery

10 All animals were housed under standard laboratory conditions, and the surgical procedures were performed in agreement with the German law (in congruence with 86/609/EEC) for the use of laboratory animals. All efforts were made to minimize the number of animals used, and all surgical procedures were performed under sufficient anesthesia to minimize animal suffering. The experimental procedures are described in detail in Bräuer et al (2001) FASEB J.
15 15:2689-2701.

Subtraction cDNA library and differential screening

The SMART cDNA technology from Clontech was used to generate high yields of full-length, double-stranded cDNA from adult, control and lesioned hippocampus rat RNA. To
20 develop the subtraction library, the Clontech PCR-Select cDNA subtraction Kit (Heidelberg, Germany) was used. Each clone was dotted in duplicates on Hybond N filters (Amersham, Germany) and screened with randomly radioactively labeled cDNA (Prime-A-Gene, Promega, Germany) from adult non-lesioned and lesioned hippocampus.

2. Analysis of the sequence of PRG-1

25 Similar to other members of LPP-family a hydrophobicity analysis of PRG-1 predicts 6 N-terminal membrane-spanning regions with a highly conserved phosphatase domain. The analysis was done using the DNAsis for Windows Version 2.6; Hitachi Software Engineering Co. Hydrophobicity Analysis Submenu using the Kyte & Doolittle algorithm with all set-
30 tings set to default values. However, unlike the other members of this family the second type of the protein consists of a long hydrophilic domain of around 400 amino acids (Fig. 2). According to the structural models of LPP orientation in the membrane, this C-terminal extension is positioned on the cytoplasmatic site and might thus play a role as a regulatory or signal transduction domain. Beside the homology of the N-terminal part of the PRG-1 to other

members of the LPP-family such as LPP-1 and the Drosophila cell migration modulator Wunen, GenBank searches revealed only one other related gene (genomic DNA sequence: GenBank acc. # NP_011255.11) for which we cloned the complete cDNA sequence and named it PRG-2. This gene shares the same C-terminal extension with partial sequence homology. Thus, these genes represent a novel distinct subclass of the LPP-1 family. Amino acid residues which have been shown to be essential for ecto-enzyme activity in the LPP-1 class of proteins are conserved in PRG-1 N-terminal sequences (Fig. 1). Database analysis of the C-terminal domains did not detect any significant similarities to any other protein or any other matches with known conserved domains (using ProDom and Swiss-Prot databases). A GenBank search for orthologous proteins showed that both genes are highly conserved in mammals (human/mouse > 93%), and partial EST sequences indicate orthologous proteins in Xenopus and Zebrafish, whereas no significant homology for the C-terminal part could be found in the Drosophila or other invertebrate genome.

3. Northern blot analysis of PRG-1 expression

The RNA isolation was described before 20 µg of total RNA from six adult control and six 1 dal animals were loaded on a 1% agarose gel containing formaldehyde, transferred to HybondTM – N s (Amersham Life Science, UK) and crosslinked by ultraviolet irradiation. As a probe for PRG-1, the full length cDNA clone as well as the C-terminal coding region was used. As probe for β-actin (as control for mRNA integrity and amounts of mRNA loaded), a cDNA fragment amplified by RT-PCR was used. Primer for the amplification of the control gene β-actin were: β-actin 5' (5' – CAC CAC AGC TGA GAG GGA AAT CGT GCG TGA – 3', SEQ ID No. 11) spanning bases 2395 – 2424, and β-actin 3'-primer (5' – ATT TGC GGT GCA GCA TGG AGG GGC CGG ACT – 3', SEQ ID no. 12) complementary to bases 3095 – 3124, with an amplicate length of 520 bp for rat β-actin cDNA (GenBank accession no. J00691). PCR was performed in 25 µl final volume containing 1 mmol/l dNTPs (Pharmacia Biotech, Germany), 2.5 units Taq Polymerase (Perkin Elmer, USA) 2.5 µl 10 x buffer including 2.5 mol/l MgCl₂ (Perkin Elmer, USA), 10 µmol/l each primer and 1 µl cDNA using a Thermo-Cycler PTC-100 (MJ Research, Inc., USA). The cycle program was: 95°C, 2 min; 35 x [94°C, 30 s; 70°C, 30 s 72°C, 2 min] and 10 min, 72°C. Both probes were labeled with the Prime-a-Gene Labelling System (Promega, USA) and [³²P] dCTP (DuPont NEN, USA). Hybridization was performed in 10 ml hybridization solution (250 mmol/l sodium-phosphate, pH 7.2, 7% SDS, 0.5 mmol/l EDTA, 1% BSA) at 60°C for 12 h. The membrane was washed in 2 x SSC at RT, 0.2 x SSC at RT, and 0.2 x SSC at 40°C for 30 min each. Membranes were ex-

posed to Kodak X-OMAT AR X-ray films at -80°C for 12 h, using an intensifying screen. Northern blot analysis revealed one distinct band which migrated around 5,5 kb. Expression of this mRNA was CNS-specific with the exception of a weak expression in testies (see Fig. 3). Thus, PRG-1 is a novel vertebrate specific protein selectively located in the brain with putative phosphatase function.

4. In situ hybridization analysis of PRG-1 expression

For hybridization, an antisense oligonucleotide (5'- GCA GAG GTC TGA ATT CTA GTG TCT ATC GTT ATA GTT CCT TAA CAG TGT GGG -3', SEQ ID No. 13) complementary to bases 425 – 475 of a rat EST cDNA clone (GenBank acc. AW 526088.1 was used). The oligonucleotide was synthesized by Metabion (Munich, Germany). The specificity was confirmed by a BLAST GenBank search to rule out cross-hybridization to other genes. The protocol was used as described by Bräuer et al (2002) *supra*.

The *in situ* hybridization analysis highlighted the tight regulation of PRG-1 transcription in the developing hippocampus. At embryonic day 16 (E16), no PRG-1 transcripts could be detected in the brain (Fig. 4, panel B). An expression signal first appears at day 19 (E19) in the subventricular zone and specifically in the hippocampal anlage, whereas other cortical regions did not show PRG-1 expression (Fig. 4, panel D). From postnatal stages on, PRG-1 mRNA is present in the hippocampus and in the entorhinal cortex throughout adult stages (see Fig. 4, panels E-J). In the dentate gyrus, a region bearing postnatally developing granule cells, weak PRG-1 mRNA expression is found in the infrapyramidal blade at P0, whereas the later developing suprapyramidal blade first showed expression signals at P5. This expression pattern remains unchanged during maturation, however, a reduced expression is apparent in the adult brain.

5. PRG-1 mRNA expression after entorhinal cortex lesion

The treatment of the animals, the surgery the construction of the subtraction cDNA library and the screening were carried out as described in example 1. The *in situ* hybridization was carried out as described in example 4. PRG-1 is upregulated one day after lesion (dal) and peaks at 5 dal in the ipsilateral hippocampus (gcl = 37%, hilus 300%, CA1 = 100%, CA3 = 60%). The contralateral hippocampus (maximum by 1 dal, gcl = 16%, hilus = 200%, CA1 = 59%, CA3 = 46%), as well as the ipsilateral cortex, shows a strong upregulation of PRG-1 mRNA (maximum by 1 dal 83%) (see Fig. 5).

6. Transfection of a PRG-1-eGFP construct

5

Antibody generation and immunohistochemistry

To design a peptide antibody against PRG-1, a sequence in the hydrophilic C-terminal region was used. The peptide (NH₂-CVGVNGDHHVPGNQ-COOH, SEQ ID No. 14), representing amino acids 490 – 507 of the PRG-1 rat sequence (GenBank acc. SEQ ID No. 5), was synthesized by BioGenes (Berlin, Germany). The amino-terminal cysteinyl residue, which is not part of the PRG-1 sequence, was included for conjugation of the peptide to a carrier protein. The peptide was conjugated through the cysteinyl sulfhydryl to maleimide activation (keyhole limpet hemocyanin). Rabbits were immunized by BioGenes. The specificity of the peptide antibody was further tested on Western blot and on brain sections by blocking via peptide incubation prior to adding the antiserum. The protocol for the immunohistochemistry is described in detail by Bräuer et al (2001) Neuroscience 102:515-526.

10
15

Western Blot analysis

For Western Blot analysis, rat adult and 5 dal hippocampus extracts were separated on a 12% SDS/PAGE and electroblotted to nitrocellulose membranes (Millipore, Germany). All incubation was done overnight at 4°C in PBST. The PRG-1 antiserum was used at a 1:2000 dilution. Secondary anti-rabbit antibody coupled with horseradish peroxidase was used at a 1:5000 dilution, and visualized by incubation in ECL detection reagents (Amersham Pharmacia, Germany). The protocol for the immunocytochemistry is essentially the same as described in detail in Bräuer et al. (2001) supra].

20
25

The immunoblot in Fig. 6D showed a single band obtained by incubation with the anti-PRG-1 antiserum. The absence of specific signal in the preimmune serum prior to immunization is noteworthy. Immunoblots from total protein extracts of adult control and deafferented hippocampus shown in Fig. 7C demonstrated an increase five days after lesion (5dal). Data represents three separate experiments in each group. Statistical difference is marked with an asterisk (mean \pm S.D.), * $P < 0.05$; Mann-Whitney-*U*-test.

30

Subcellular localization

PRG-1 tagged with the eGFP reporter gene was used to identify the subcellular localization. Golgi apparatus was visualized with the cell tracker BODYPY ceramide (Molecular Probes, Netherlands). The staining protocol was obtained from molecular Probes.

5 The transfection studies using a PRG-1 construct tagged with a eGFP reporter gene revealed that PRG-1 protein was processed in COS-7 cells through the Golgi apparatus (data not shown) to its final localization in the plasma membrane of small processes (Fig. 6, panel A). To localize the PRG-1 proteins *in vivo* an antiserum against a peptide from the cytoplasmic C-terminus of PRG-1 was raised. This antiserum specifically stained transfected COS-7 cells,
10 which expressed PRG-1-eGFP fusion proteins (Fig. 6, panel A-C) and detected a specific band in Western blot analysis (Fig. 6, panel B). Both the immunostaining and Western blot signal could be blocked by specific peptide incubation prior to the addition of antiserum (data not shown). The fluorescence of the labeled entire-PRG-1 peptide antibody and the fluorescence of the eGFP part of the fusion protein colocalized in COS-7 cells and in the processes
15 (see white arrows in panels A-C of Fig. 6).

7. Expression analysis by RT-PCR

Tissue from the retraction assays, E16 and P0 explants and N1E-115 cells were used for mRNA isolation and as templates for RT-PCR. cDNA from testis was used as a positive control. The MidiMACS mRNA Isolation Kit (Miltenyi Biotec, Germany) was used to isolate
20 mRNA from the explants or cells. Reverse transcription was performed as described by Bräuer et al. (2000) Hippocampus 10:632-644. PCR was performed with the following primers: PRG-1-5'-primer (5'-CTA GGC TTG TAG CTG TGG GGA ATT TC-3', SEQ ID No. 15), spanning bases 896 bp –921 bp, and PRG-1-3'-primer (5'-TCA ATC CTT ATA AGC
25 CCG TGT G-3', SEQ ID No. 16) complementary to bases 2202 bp – 2225 bp with an amplification length of 1329 bp of the PRG-1 cDNA (SEQ ID No. 10). For amplification of the EDG receptor cDNA, the primer EDG-2-5' primer (5'-GAA CTT TGC GAG TGA GCT GG-3', SEQ ID No. 17), spanning bases 836 bp –855 bp, and EDG-2-3'-primer (5'TGC GGA GAG CTT TAA CCT CC-3', SEQ ID NO. 18), complementary to bases 1165 bp – 1184 bp
30 with an amplification length of 348 bp of the EDG-2 cDNA (GenBank accession no. NM010336), the primer EDG-4-5'-primer (5'-CCT ACC TCT TCC TCA TGT TC-3', SEQ ID No. 19), spanning bases 344 bp – 363 bp, and EDG-4-3'-primer (5'TAA AGG GTG GAG TCC ATC AG-3', SEQ ID No. 20), complementary to bases 1199 bp – 1148 bp of the EDG-4 cDNA (GenBank accession no. NM020028), the primer EDG-7-5'-primer (5'-GGA ATT GCC

TCT GCA ACA TCT-3', SEQ ID No. 21), spanning bases 673 bp – 693 bp, and EDG-7-3' primer (5'-GAG TAG ATG ATG GG TTC A-3', SEQ ID No. 22), complementary to bases 1096 bp – 1054 bp of the EDG-7 cDNA (GenBank accession no. NM022983). PCR was performed using a Thermo-cycler PTC-100 (MJ Research, Inc.) in 25 µl final volume containing 10 µmol/l dNTPs (Pharmacia, Germany), 2.5 units Taq Polymerase (Stratagen, Germany), 2.5 µl 10 x buffer including 2.5 mol/l MgCl₂ (Stratagen, Germany), 10 µmol/l of each primer, and 2 µl of each cDNA for all molecular analysis. For all EDG receptors, as well as for PRG-1 amplification, the cycle program was: 2 min at 95°C, 40 x (94°C, 30 sec; 52°C, 30 sec; and 72°C, 1 min), and 5 min at 72°C. Amplification of β-actin cDNA was performed as described by Bräuer et al. (2000) Hippocampus 10:632-644.

8. Immunocytochemical staining of rat hippocampus

The antibody described in experiment 6 was also used for the immunocytochemical staining and Western blot analysis. For Western blot analysis, rat adult and 5dal hippocampus extracts were separated on a 12% SDS/PAGE and electroblotted to nitrocellulose membranes (Millipore, Germany). All incubation was done overnight at 4°C in PBST. The PRG-1 antiserum was used at a 1:2000 dilution. Secondary anti-rabbit antibody coupled with horseradish peroxidase was used at a 1:5000 dilution, and visualized by incubation in ECL detection reagents (Amersham Pharmacia, Germany). The protocol for the immunocytochemistry is essentially the same as described in detail in Bräuer et al., 2001 (FASEB J, 15, 2689-2701). The immunocytochemical staining of rat hippocampus revealed that PRG-1 was specifically expressed in neurons (see Fig. 7, panel A) and in particular pyramidal neurons are labeled in the CA1 and CA3 region, polymorphic cells are stained in the hilus and granule cells of the dentate gyrus are also immunopositive. However, the outer molecular layer, the termination zone of afferents from the entorhinal cortex (Skutella T. and Nitsch R. (2001) Trends. Neuroscience 24:107-163) showed no PRG-1 positive fibers (Fig. 7, panel A). Conversely five days after entorhinal cortex lesion, a clear immunoreactive PRG-1 positive band appeared in the denervated outer molecular layer (Fig. 7, panel B). Western blot analysis revealed a 50% increase in PRG-1 expression in the denervated hippocampus (Fig. 7, panel C). PRG-1 immunostaining highlighted single axonal processes in the outer molecular layer which form terminal branches (Fig. 8, panel A). The higher magnification of an area from the immuno stained axons showed that PRG-1 is indeed localized in the growth cone-like axon structures in the denervated zones of the hippocampus (see Fig. 8, panel B).

Subcellular localization

- 5 PRG-1 tagged with the eGFP reporter gene was used to identify the subcellular localization. Golgi apparatus was visualized with the cell tracker BODYPY ceramide (Molecular Probes, Oregon). The staining protocol was obtained from Molecular Probes.

9. Effect of PRG-1 expression on the LPA response of neurons

10

LPA induced neurite retraction in explants

- Entorhinal explants from E16 and P0 rat pups were obtained from timed-pregnant Wistar rats and were cultivated as described (N. E. Savaskan (2000) Eur. J. Neurosci 12:1024-1032). In brief, entorhinal cortex was carefully dissected from the hippocampal anlage and the meninges were removed. Explants were gently transferred with a fire-polished Pasteur pipette into 12-well plates and cultivated on baked glass cover slides coated with laminin and poly-L-lysine (25 µg/ml and 10 µg/ml, respectively) in culture medium containing selenium-defined fetal bovine serum [5%] (N. E. Savaskan et al (2000) FASEB J. e-published) (Neurobasal medium plus 25 µg/ml Penicillin/Streptomycin; B-27 supplement). After 24 h, culture medium was exchanged and cultivation was further performed in serum-free Neurobasal medium for 20 h. Serum-starved explants were treated with 100 nmol/l oleoyl-LPA (5 mmol/l stock solution in ultra-filtrated water) for 10 min or with vehicle (0.9% NaCl) and then fixed in 4% paraformaldehyde for 20 min. For F-actin staining, fixed tissues were incubated with TRITC-phalloidin (0.1 µg/ml, Sigma, Germany) for 40 min, followed by incubation with HOECHST 33258 dye (1:20,000, Sigma, Germany) for 5 min at room temperature. After three washing steps in PBS, explants were coverslipped with ImmunoMount (Merck, Germany) prior to analysis. Images were taken with a CCD camera on an Olympus BX-50 microscope and quantification was performed using the Meta Morph analysis system (Universal Imaging, PA). For statistical analysis Statview II was used (Abacus, USA).

30

A dose response of LPA neurite length of E16 and P0 explants was done as described above, however, with 0, 0.1, 1.0 and 10 µmol/l.

Entorhinal explants obtained at day 16 (E16) do not express PRG-1 while postnatal explants (P0) express PRG-1 (see Fig. 9, panel A). In contrast, the LPA-specific receptors EDG-2/4/7 were equally expressed in both embryonic and postnatal explants (see Fig. 9B). The control RT-PCR with β -actin showed the integrity of the RNA from E16 and P0 cells. Both embryonic and postnatal explants grow equally well under serum-free culture conditions and show long extending axons (see Fig. 9, panels D and F). However, their response to LPA differed dramatically (see Fig. 9, panels E and G). Whereas application of 100 nmol/l LPA led to rapid neurite retraction in embryonic entorhinal explants (E16; n = 20, compare panel D to panel E), postnatal explants (P0; n = 22, compare panel F to panel G) did not differ significantly from vehicle treated control cultures. Thus, postnatal entorhinal axons expressing PRG-1 are resistant to LPA-induced neurite retraction. The amount of retraction observed in panels D-G is quantified in Fig. 10, panel A. The dose response is shown in Fig. 10, panel B.

10. Differential effect of PRG-1 and PRG-1 mutant on LPA-induced neurite retraction

Site-directed mutagenesis of PRG-1^{HIS/LYS}

The rat PRG-1 full length clone was amplified by Marathon PCR (Clontech, USA) from adult rat hippocampus RNA (SEQ ID No. 10). For transfection studies, the full length PRG-1 coding sequence was fused to EGFP (pEGFP-N1 vector Clontech, USA). The PRG-1^{His/Lys} exchange mutant at the catalytic histidine (His-252) was introduced in the same protein fusion vector by site specific mutagenesis (CAT to AAG).

LPA induced neurite retraction and protection in N1E-115 cells

N1E-115 mouse neuroblastoma cells (ATCC: CRL-2263) were routinely grown in DMEM medium supplemented with selenium-defined fetal bovine serum (10%). The cells were seeded on baked glass cover slides at a density of 10,000 cells/cm². The next day, cells were transfected with the cationic lipids procedure (FuGene6, Roche, Germany) and cultivated for 24 h. Serum-starvation was performed for 20 h in DMEM medium, followed by treatment with 10 μ mol/l oleoyl-LPA or vehicle (0.9% sodium chloride) (K. Jalink et al (1993) Cell Growth Differ. 4:247-255). After 10 min, cells were fixed in 4% paraformaldehyde for 20 min at room temperature and further analysis processed as described above. Images were taken with a CCD camera on an Olympus BX-50 microscope. Quantification was performed with the Meta Morph analysis system (University Imaging, PA). For statistical analysis Stat-view II was used (Abacus, USA). The effects of transfection on cell viability were analysed

by MTT assay and propidium iodide staining (Savaskan N.E. et al. (2002) FASEB J. 17:112-114).

N1E-115 cells are uniformly sensitive to LPA-induced growth cone collapse (see Fig. 11 and compare first row of panels versus second row of panels). This is also confirmed by phalloidin staining (K. Jalink et al, (1994) J. Cell Biol. 126:801-810) which showed actin polymerization upon PRG-1 overexpression, however, led to a resistance of N1E-115 cells to LPA-induced growth cone collapse and also prevented LPA-induced actin-polymerization (see Fig. 11, third row of panels). The mutation of the conserved catalytic histidine (His-252) to a lysine (PRG-1^{His/Lys}) a change which has been shown to completely abolish enzymatic function of the catalytic center of LPP-1 (N. Zhang, et al. (1997) *supra*) no longer prevented LPA-induced retraction of processes as achieved by the wt-construct. This shows that the conserved enzymatic domain of the LPP-1 family is necessary for PRG-1 function in attenuating LPA-induced neurite retraction. The results of the experiments are quantified in Fig. 12, panel A. The transfection, however, had no effect on the expression of the LPA-receptors EDG-2/4/7 (see Fig. 12, panel B). The control for the integrity of the RNA tested by RT-PCR is shown in Fig. 12, panel C. The resistance against CPA-induced growth cone collapse achieved by PRG-1 overexpression could only be overcome by a 10-fold increase of LPA applied to the culture (Fig. 12, panel D).

11. Ectophosphatase activity in PRG-1 transfected N1H-115

The assay procedures used were adapted from those described in Imai et al. (2000) J. Clin. Endocrinol. Metab 85:3370-3375 and Hooks S.B. et al. (2001) J. Biol. Chem. 276:4611-4621. Briefly, transfected cells were harvested after serum-free cultivation for 20 h by scraping in lysis buffer (containing 20 mmol/l HEPES pH 7.4, 1 mmol/l NaHCO₃, 500 µmol/l DTT, 1 mmol/l EGTA) and sonicated with two strokes (Bandolin GM 70, Germany). After centrifugation at 800 x g, the supernatant was diluted in lysis buffer and centrifuged at 100,000 x g for 1 h. Alternatively, the supernatant was topped on a sucrose step gradient (50% and 5%) and centrifuged at 100,000 x g for 1 h (Savaskan, N.E. et al. (2000) Eur. J. Neurosci. 12:1024-1032. The resulting pellet and interphase, respectively, were re-homogenized and centrifuged again at 100,000 x g for 1 h. The crude membrane pellet was re-suspended in reaction buffer (50 mmol/l HEPES pH 7.5; 1 mmol/l EGTA) and protein concentration was determined spectroscopically by Bradford protein assay (Amersham, Germany). Assessment of LPA metabolism was performed with exogenous ³H-oleoeyl-LPA (Perkin Elmer, Germany) measur-

ing ^3H -oleoeyl-glycerol production. Briefly, 10-25 μg membrane proteins were pre-warmed to 37°C in reaction buffer and reactions were started by the addition of 10 $\mu\text{mol/l}$ ^3H -oleoeyl-LPA. The reaction was allowed to proceed for 5-30 min at 37°C and stopped with the addition of 2.5 vol acidified methanol and 1.5 vol chloroform. After two-phase separation, the chloroform phase was dried under N_2 and applied to silica gel matrices (Machery Nagel, Germany). Plates were developed in chloroform-aceton-acetic acid (90:10:1). All fractions of dried plates were scintillation counted and compared with authentic standards (LPA, 1-monoolein, 1,2-monoolein, oleic acid). All experiments were performed in triplicate. The ecto-phosphatase assay in intact cells was determined as described above. ^3H -LPA was added to the serum-free medium and the reaction was allowed to proceed for 5-60 min. The reaction was stopped by adding 2.5 vol acidified methanol and 1.5 vol chloroform to the supernatant. The radioactivity was determined in the lipid fraction as described above.

The results from three independent sets of experiments are shown in Fig. 12, panel F (one set with $n = 40$ for each group in the outgrowth assay). PRG-1-eGFP showed essentially the same ecto-phosphatase activity as the wt PRG-1 construct. Statistical differences from controls are marked with an asterisk (mean \pm S.D.), *** $P < 0.001$; two-tailed t test with Bonferroni correction for multiple comparisons.

The extracellular LPA-degradation achieved by transfected N1E-115 cells revealed a 5-fold increase in ecto-phosphatase activity of wildtype PRG-1 transfectants when compared to eGFP transfection alone (Fig. 12, panel F). Moreover, point-mutation in the conserved catalytic domain by His/Lys exchange led to a 95 % decrease of ecto-phosphatase activity (Fig. 12, panel F). These findings show that PRG-1 has ecto-phosphatase activity which is conveyed by a conserved enzymatic domain present in the LPP-1 family, and necessary for attenuating LPA-induced neurite retraction. In addition transfection of the PRG-1 construct into N1E-115 cells did not prevent CPA-induced retraction of processes as achieved by the wt-constructs, whereas cell viability was unaffected (see Fig. 12, panel E).

Thus the ecto-phosphatase activity of PRG-1 was shown by two independent experiments: a) transfection of a mutant construct into N1E-115 cells did not prevent LPA-induced retraction of processes as achieved by transfection of wt construct and b) the transfection of the wt construct led to a 5-fold increase in ecto-phosphatase activity, if compared to cells transfected with a control vector.

12. Model of axon growth mechanism

- 5 Fig. 13 shows a diagram of the proposed axon growth mechanism in a phospholipid rich environment. Axons that are sensitive to repulsive phosphor lipid but do not express PRG-1 are unable to cross a phosphor lipid-rich barrier. In contrasts PRG-1 expressing neurons can grow through a phosphor lipid rich zone by local depleting the extracellular pool of repulsive phosphor lipids acting as ligands on EDG-receptors. This way, PRG-1 may regulate the activation
- 10 of EDG-receptors and thereby modulate axonal outgrowth.

THIS PAGE BLANK (USPTO)

EPO - Munich
83
11 Feb. 2003

SEQUENCE LISTING

<110> Universitätsklinikum Charité

<120> Lipid phosphate phosphatases and uses thereof for treating neuronal diseases

<130> FB11828

<150> EP 02 020 679.3

<151> 2002-09-13

<160> 22

<170> PatentIn version 3.2

<210> 1

<211> 763

<212> PRT

<213> Homo sapiens

<400> 1

Met	Gln	Arg	Ala	Gly	Ser	Ser	Gly	Gly	Arg	Gly	Glu	Cys	Asp	Ile	Ser
1				5					10					15	

Gly	Ala	Gly	Arg	Leu	Gly	Leu	Glu	Glu	Ala	Ala	Arg	Leu	Ser	Cys	Ala
			20					25					30		

Val	His	Thr	Ser	Pro	Gly	Gly	Gly	Arg	Arg	Pro	Gly	Gln	Ala	Ala	Gly
		35					40					45			

Met	Ser	Ala	Lys	Glu	Arg	Pro	Lys	Gly	Lys	Val	Ile	Lys	Asp	Ser	Val
	50					55					60				

Thr	Leu	Leu	Pro	Cys	Phe	Tyr	Phe	Val	Glu	Leu	Pro	Ile	Leu	Ala	Ser
65					70					75					80

Ser	Val	Val	Ser	Leu	Tyr	Phe	Leu	Glu	Leu	Thr	Asp	Val	Phe	Lys	Pro
				85					90					95	

Val	His	Ser	Gly	Phe	Ser	Cys	Tyr	Asp	Arg	Ser	Leu	Ser	Met	Pro	Tyr
			100					105					110		

Ile	Glu	Pro	Thr	Gln	Glu	Ala	Ile	Pro	Phe	Leu	Met	Leu	Leu	Ser	Leu
		115					120					125			

Ala	Phe	Ala	Gly	Pro	Ala	Ile	Thr	Ile	Met	Val	Gly	Glu	Gly	Ile	Leu
	130					135					140				

Tyr	Cys	Cys	Leu	Ser	Lys	Arg	Arg	Asn	Gly	Val	Gly	Leu	Glu	Pro	Asn
145					150					155					160

Ile Asn Ala Gly Gly Cys Asn Phe Asn Ser Phe Leu Arg Arg Ala Val
 165 170 175

Arg Phe Val Gly Val His Val Phe Gly Leu Cys Ser Thr Ala Leu Ile
 180 185 190

Thr Asp Ile Ile Gln Leu Ser Thr Gly Tyr Gln Ala Pro Tyr Phe Leu
 195 200 205

Thr Val Cys Lys Pro Asn Tyr Thr Ser Leu Asn Val Ser Cys Lys Glu
 210 215 220

Asn Ser Tyr Ile Val Glu Asp Ile Cys Ser Gly Ser Asp Leu Thr Val
 225 230 235 240

Ile Asn Ser Gly Arg Lys Ser Phe Pro Ser Gln His Ala Thr Leu Ala
 245 250 255

Ala Phe Ala Ala Val Tyr Val Ser Met Tyr Phe Asn Ser Thr Leu Thr
 260 265 270

Asp Ser Ser Lys Leu Leu Lys Pro Leu Leu Val Phe Thr Phe Ile Ile
 275 280 285

Cys Gly Ile Ile Cys Gly Leu Thr Arg Ile Thr Gln Tyr Lys Asn His
 290 295 300

Pro Val Asp Val Tyr Cys Gly Phe Leu Ile Gly Gly Gly Ile Ala Leu
 305 310 315 320

Tyr Leu Gly Leu Tyr Ala Val Gly Asn Phe Leu Pro Ser Asp Glu Ser
 325 330 335

Met Phe Gln His Arg Asp Ala Leu Arg Ser Leu Thr Asp Leu Asn Gln
 340 345 350

Asp Pro Asn Arg Leu Leu Ser Ala Lys Asn Gly Ser Ser Ser Asp Gly
 355 360 365

Ile Ala His Thr Glu Gly Ile Leu Asn Arg Asn His Arg Asp Ala Ser
 370 375 380

Ser Leu Thr Asn Leu Lys Arg Ala Asn Ala Asp Val Glu Ile Ile Thr
 385 390 395 400

Pro Arg Ser Pro Met Gly Lys Glu Asn Met Val Thr Phe Ser Asn Thr
 405 410 415

Leu Pro Arg Ala Asn Thr Pro Ser Val Glu Asp Pro Val Arg Arg Asn
 420 425 430

Ala Ser Ile His Ala Ser Met Asp Ser Ala Arg Ser Lys Gln Leu Leu
 435 440 445

Thr Gln Trp Lys Asn Lys Asn Glu Ser Arg Lys Leu Ser Leu Gln Val
 450 455 460

Ile Glu Pro Glu Pro Gly Gln Ser Pro Pro Arg Ser Ile Glu Met Arg
 465 470 475 480

Ser Ser Ser Glu Pro Ser Arg Val Gly Val Asn Gly Asp His His Gly
 485 490 495

Pro Gly Asn Gln Tyr Leu Lys Ile Gln Pro Gly Ala Val Pro Gly Cys
 500 505 510

Asn Asn Ser Met Pro Gly Gly Pro Arg Val Ser Ile Gln Ser Arg Pro
 515 520 525

Gly Ser Ser Gln Leu Val His Ile Pro Glu Glu Thr Gln Glu Asn Ile
 530 535 540

Ser Thr Ser Pro Lys Ser Ser Ser Ala Arg Ala Lys Trp Leu Lys Ala
 545 550 555 560

Ala Glu Lys Thr Val Ala Cys Asn Arg Ser Asn Ser Gln Pro Arg Ile
 565 570 575

Met Gln Val Ile Ala Met Ser Lys Gln Gln Gly Val Leu Gln Ser Ser
 580 585 590

Pro Lys Asn Thr Glu Gly Ser Thr Val Ser Cys Thr Gly Ser Ile Arg
 595 600 605

Tyr Lys Thr Leu Thr Asp His Glu Pro Ser Gly Ile Val Arg Val Glu
 610 615 620

Ala His Pro Glu Asn Asn Arg Pro Ile Ile Gln Ile Pro Ser Thr Glu
 625 630 635 640

Gly Glu Gly Ser Gly Ser Trp Lys Trp Lys Ala Pro Glu Lys Gly Ser
 645 650 655

Leu Arg Gln Thr Tyr Glu Leu Asn Asp Leu Asn Arg Asp Ser Glu Ser
 660 665 670

Cys Glu Ser Leu Lys Asp Ser Phe Gly Ser Gly Asp Arg Lys Arg Ser
675 680 685

Asn Ile Asp Ser Asn Glu His His His His Gly Ile Thr Thr Ile Arg
690 695 700

Val Thr Pro Val Glu Gly Ser Glu Ile Gly Ser Glu Thr Leu Ser Ile
705 710 715 720

Ser Ser Ser Arg Asp Ser Thr Leu Arg Arg Lys Gly Asn Ile Ile Leu
725 730 735

Ile Pro Glu Arg Ser Asn Ser Pro Glu Asn Thr Arg Asn Ile Phe Tyr
740 745 750

Lys Gly Thr Ser Pro Thr Arg Ala Tyr Lys Asp
755 760

<210> 2
<211> 746
<212> PRT
<213> Homo sapiens

<400> 2

Met Ile Ser Thr Lys Glu Lys Asn Lys Ile Pro Lys Asp Ser Met Thr
1 5 10 15

Leu Leu Pro Cys Phe Tyr Phe Val Glu Leu Pro Ile Val Ala Ser Ser
20 25 30

Ile Val Ser Leu Tyr Phe Leu Glu Leu Thr Asp Leu Phe Lys Pro Ala
35 40 45

Lys Val Gly Phe Gln Cys Tyr Asp Arg Thr Leu Ser Met Pro Tyr Val
50 55 60

Glu Thr Asn Glu Glu Leu Ile Pro Leu Leu Met Leu Leu Ser Leu Ala
65 70 75 80

Phe Ala Ala Pro Ala Ala Ser Ile Met Val Ala Glu Gly Met Leu Tyr
85 90 95

Cys Leu Gln Ser Arg Leu Trp Gly Arg Ala Gly Gly Pro Ala Gly Ala
100 105 110

Glu Gly Ser Ile Asn Ala Gly Gly Cys Asn Phe Asn Ser Phe Leu Arg
115 120 125

Arg Thr Val Arg Phe Val Gly Val His Val Phe Gly Leu Cys Ala Thr
 130 135 140

Ala Leu Val Thr Asp Val Ile Gln Leu Ala Thr Gly Tyr His Thr Pro
 145 150 155 160

Phe Phe Leu Thr Val Cys Lys Pro Asn Tyr Thr Leu Leu Gly Thr Ser
 165 170 175

Cys Glu Val Asn Pro Tyr Ile Thr Gln Asp Ile Cys Ser Gly His Asp
 180 185 190

Ile His Ala Ile Leu Ser Ala Arg Lys Thr Phe Pro Ser Gln His Ala
 195 200 205

Thr Leu Ser Ala Phe Ala Ala Val Tyr Val Ser Val Ser Pro Ala Pro
 210 215 220

His Cys Pro Ser Gln Ala Leu Leu Leu Thr Arg Gly Glu Pro Ser Leu
 225 230 235 240

Thr Pro Thr Pro Met Pro Gln Met Tyr Phe Asn Ser Val Ile Ser Asp
 245 250 255

Thr Thr Lys Leu Leu Lys Pro Ile Leu Val Phe Ala Phe Ala Ile Ala
 260 265 270

Ala Gly Val Cys Gly Leu Thr Gln Ile Thr Gln Tyr Arg Ser His Pro
 275 280 285

Val Asp Val Tyr Ala Gly Phe Leu Ile Gly Ala Gly Ile Ala Ala Tyr
 290 295 300

Leu Ala Cys His Ala Val Gly Asn Phe Gln Ala Pro Pro Ala Glu Lys
 305 310 315 320

Pro Ala Ala Pro Ala Pro Ala Lys Asp Ala Leu Arg Ala Leu Thr Gln
 325 330 335

Arg Gly His Asp Ser Val Tyr Gln Gln Asn Lys Ser Val Ser Thr Asp
 340 345 350

Glu Leu Gly Pro Pro Gly Arg Leu Glu Gly Ala Pro Arg Pro Val Ala
 355 360 365

Arg Glu Lys Thr Ser Leu Gly Ser Leu Lys Arg Ala Ser Val Asp Val

370	375	380
Asp Leu Leu Ala Pro Arg Ser Pro Met Ala Lys Glu Asn Met Val Thr 385	390	395 400
Phe Ser His Thr Leu Pro Arg Ala Ser Ala Pro Ser Leu Asp Asp Pro 405	410	415
Ala Arg Arg His Met Thr Ile His Val Pro Leu Asp Ala Ser Arg Ser 420	425	430
Lys Gln Leu Ile Ser Glu Trp Lys Gln Lys Ser Leu Glu Gly Arg Gly 435	440	445
Leu Gly Leu Pro Asp Asp Ala Ser Pro Gly His Leu Arg Ala Pro Ala 450	455	460
Glu Pro Met Ala Glu Glu Glu Glu Glu Glu Asp Glu Glu Glu Glu 465	470	475 480
Glu Glu Glu Glu Glu Glu Glu Asp Glu Gly Pro Ala Pro Pro Ser Leu 485	490	495
Tyr Pro Thr Val Gln Ala Arg Pro Gly Leu Gly Pro Arg Val Ile Leu 500	505	510
Pro Pro Arg Ala Gly Pro Pro Pro Leu Val His Ile Pro Glu Glu Gly 515	520	525
Ala Gln Thr Gly Ala Gly Leu Ser Pro Lys Ser Gly Ala Gly Val Arg 530	535	540
Ala Lys Trp Leu Met Met Ala Glu Lys Ser Gly Ala Ala Val Ala Asn 545	550	555 560
Pro Pro Arg Leu Leu Gln Val Ile Ala Met Ser Lys Ala Pro Gly Ala 565	570	575
Pro Gly Pro Lys Ala Ala Glu Thr Ala Ser Ser Ser Ser Ala Ser Ser 580	585	590
Asp Ser Ser Gln Tyr Arg Ser Pro Ser Asp Arg Asp Ser Ala Ser Ile 595	600	605
Val Thr Ile Asp Ala His Ala Pro His His Pro Val Val His Leu Ser 610	615	620

Ala Gly Gly Ala Pro Trp Glu Trp Lys Ala Ala Gly Gly Gly Ala Lys
625 630 635 640

Ala Glu Ala Asp Gly Gly Tyr Glu Leu Gly Asp Leu Ala Arg Gly Phe
645 650 655

Arg Gly Gly Ala Lys Pro Pro Gly Val Ser Pro Gly Ser Ser Val Ser
660 665 670

Asp Val Asp Gln Glu Glu Pro Arg Phe Gly Ala Val Ala Thr Val Asn
675 680 685

Leu Ala Thr Gly Glu Gly Leu Pro Pro Leu Gly Ala Ala Asp Gly Ala
690 695 700

Leu Gly Pro Gly Ser Arg Glu Ser Thr Leu Arg Arg His Ala Gly Gly
705 710 715 720

Leu Gly Leu Ala Glu Arg Glu Ala Glu Ala Glu Ala Glu Gly Tyr Phe
725 730 735

Arg Lys Met Gln Ala Arg Arg Phe Pro Asp
740 745

<210> 3
<211> 766
<212> PRT
<213> Mus musculus

<400> 3

Met Gln Arg Ala Gly Ser Ser Gly Ala Arg Gly Glu Cys Asp Ile Ser
1 5 10 15

Gly Ala Gly Arg Leu Arg Leu Glu Gln Ala Ala Arg Leu Gly Gly Arg
20 25 30

Thr Val His Thr Ser Pro Gly Gly Gly Leu Gly Ala Arg Gln Ala Ala
35 40 45

Gly Met Ser Ala Lys Glu Arg Pro Lys Gly Lys Val Ile Lys Asp Ser
50 55 60

Val Thr Leu Leu Pro Cys Phe Tyr Phe Val Glu Leu Pro Ile Leu Ala
65 70 75 80

Ser Ser Val Val Ser Leu Tyr Phe Leu Glu Leu Thr Asp Val Phe Lys
85 90 95

Pro Val His Ser Gly Phe Ser Cys Tyr Asp Arg Ser Leu Ser Met Pro
 100 105 110

Tyr Ile Glu Pro Thr Gln Glu Ala Ile Pro Phe Leu Met Leu Leu Ser
 115 120 125

Leu Ala Phe Ala Gly Pro Ala Ile Thr Ile Met Val Gly Glu Gly Ile
 130 135 140

Leu Tyr Cys Cys Leu Ser Lys Arg Arg Asn Gly Ala Gly Leu Glu Pro
 145 150 155 160

Asn Ile Asn Ala Gly Gly Cys Asn Phe Asn Ser Phe Leu Arg Arg Ala
 165 170 175

Val Arg Phe Val Gly Val His Val Phe Gly Leu Cys Ser Thr Ala Leu
 180 185 190

Ile Thr Asp Ile Ile Gln Leu Ser Thr Gly Tyr Gln Ala Pro Tyr Phe
 195 200 205

Leu Thr Val Cys Lys Pro Asn Tyr Thr Ser Leu Asn Val Ser Cys Lys
 210 215 220

Glu Asn Ser Tyr Ile Val Glu Asp Ile Cys Ser Gly Ser Asp Leu Thr
 225 230 235 240

Val Ile Asn Ser Gly Arg Lys Ser Phe Pro Ser Gln His Ala Thr Leu
 245 250 255

Ala Ala Phe Ala Ala Val Tyr Val Ser Met Tyr Phe Asn Ser Thr Leu
 260 265 270

Thr Asp Ser Ser Lys Leu Leu Lys Pro Leu Leu Val Phe Thr Phe Ile
 275 280 285

Ile Cys Gly Ile Ile Cys Gly Leu Thr Arg Ile Thr Gln Tyr Lys Asn
 290 295 300

His Pro Val Asp Val Tyr Cys Gly Phe Leu Ile Gly Gly Gly Ile Ala
 305 310 315 320

Leu Tyr Leu Gly Leu Tyr Ala Val Gly Asn Phe Leu Pro Ser Glu Asp
 325 330 335

Ser Met Leu Gln His Arg Asp Ala Leu Arg Ser Leu Thr Asp Leu Asn
 340 345 350

Gln Asp Pro Ser Arg Val Leu Ser Ala Lys Asn Gly Ser Ser Gly Asp
 355 360 365

Gly Ile Ala His Thr Glu Gly Ile Leu Asn Arg Asn His Arg Asp Ala
 370 375 380

Ser Ser Leu Thr Asn Leu Lys Arg Ala Asn Ala Asp Val Glu Ile Ile
 385 390 395 400

Thr Pro Arg Ser Pro Met Gly Lys Glu Ser Met Val Thr Phe Ser Asn
 405 410 415

Thr Leu Pro Arg Ala Asn Thr Pro Ser Val Glu Asp Pro Val Arg Arg
 420 425 430

Asn Ala Ser Ile His Ala Ser Met Asp Ser Ala Arg Ser Lys Gln Leu
 435 440 445

Leu Thr Gln Trp Lys Ser Lys Asn Glu Ser Arg Lys Met Ser Leu Gln
 450 455 460

Val Met Asp Thr Glu Pro Glu Gly Gln Ser Pro Pro Arg Ser Ile Glu
 465 470 475 480

Met Arg Ser Ser Ser Glu Pro Ser Arg Val Gly Val Asn Gly Asp His
 485 490 495

His Val Pro Gly Asn Gln Tyr Leu Lys Ile Gln Pro Gly Thr Val Pro
 500 505 510

Gly Cys Asn Asn Ser Met Pro Gly Gly Pro Arg Val Ser Ile Gln Ser
 515 520 525

Arg Pro Gly Ser Ser Gln Leu Val His Ile Pro Glu Glu Thr Gln Glu
 530 535 540

Asn Ile Ser Thr Ser Pro Lys Ser Ser Ser Ala Arg Ala Lys Trp Leu
 545 550 555 560

Lys Ala Ala Glu Lys Thr Val Asp Cys Asn Arg Ser Asn Asn Gln Pro
 565 570 575

Arg Ile Met Gln Val Ile Ala Met Ser Lys Gln Gln Gly Val Leu Gln
 580 585 590

Ser Ser Pro Lys Asn Ala Glu Gly Ser Thr Val Thr Cys Thr Gly Ser
 595 600 605

Ile Arg Tyr Lys Thr Leu Thr Asp His Glu Pro Ser Gly Ile Val Arg
610 615 620

Val Glu Ala His Pro Glu Asn Asn Arg Pro Ile Ile Gln Ile Pro Ser
625 630 635 640

Ser Thr Glu Gly Glu Gly Ser Gly Ser Trp Lys Trp Lys Val Pro Glu
645 650 655

Lys Ser Ser Leu Arg Gln Thr Tyr Glu Leu Asn Asp Leu Asn Arg Asp
660 665 670

Ser Glu Ser Cys Glu Ser Leu Lys Asp Ser Phe Gly Ser Gly Asp Arg
675 680 685

Lys Arg Ser Asn Ile Asp Ser Asn Glu His His His His Gly Ile Thr
690 695 700

Thr Ile Arg Val Thr Pro Val Glu Gly Ser Glu Ile Gly Ser Glu Thr
705 710 715 720

Leu Ser Val Ser Ser Ser Arg Asp Ser Thr Leu Arg Arg Lys Gly Asn
725 730 735

Ile Ile Leu Ile Pro Glu Arg Ser Asn Ser Pro Glu Asn Thr Arg Asn
740 745 750

Ile Phe Tyr Lys Gly Thr Ser Pro Thr Arg Ala Tyr Lys Asp
755 760 765

<210> 4

<211> 716

<212> PRT

<213> Mus musculus

<400> 4

Met Leu Ala Met Lys Glu Lys Asn Lys Thr Pro Lys Asp Ser Met Thr
1 5 10 15

Leu Leu Pro Cys Phe Tyr Phe Val Glu Leu Pro Ile Val Ala Ser Ser
20 25 30

Ile Val Ser Leu Tyr Phe Leu Glu Leu Thr Asp Leu Phe Lys Pro Ala
35 40 45

Lys Val Gly Phe Gln Cys Tyr Asp Arg Ala Leu Ser Met Pro Tyr Val
50 55 60

Glu Thr Asn Glu Glu Leu Ile Pro Leu Leu Met Leu Leu Ser Leu Ala
65 70 75 80

Phe Ala Ala Pro Ala Ala Ser Ile Met Val Gly Glu Gly Met Val Tyr
85 90 95

Cys Leu Gln Ser Arg Leu Trp Gly Arg Gly Pro Gly Gly Val Glu Gly
100 105 110

Ser Ile Asn Ala Gly Gly Cys Asn Phe Asn Ser Phe Leu Arg Arg Thr
115 120 125

Val Arg Phe Val Gly Val His Val Phe Gly Leu Cys Ala Thr Ala Leu
130 135 140

Val Thr Asp Val Ile Gln Leu Ala Thr Gly Tyr His Thr Pro Phe Phe
145 150 155 160

Leu Thr Val Cys Lys Pro Asn Tyr Thr Leu Leu Gly Thr Ser Cys Glu
165 170 175

Ser Asn Pro Tyr Ile Thr Gln Asp Ile Cys Ser Gly His Asp Thr His
180 185 190

Ala Ile Leu Ser Ala Arg Lys Thr Phe Pro Ser Gln His Ala Thr Leu
195 200 205

Ser Ala Phe Ala Ala Val Tyr Val Ser Met Tyr Phe Asn Ala Val Ile
210 215 220

Ser Asp Thr Thr Lys Leu Leu Lys Pro Ile Leu Val Phe Ala Phe Ala
225 230 235 240

Ile Ala Ala Gly Val Cys Gly Leu Thr Gln Ile Thr Gln Tyr Arg Ser
245 250 255

His Pro Val Asp Val Tyr Ala Gly Phe Leu Ile Gly Ala Gly Ile Ala
260 265 270

Ala Tyr Leu Ala Cys His Ala Val Gly Asn Phe Gln Ala Pro Pro Ala
275 280 285

Glu Lys Val Pro Thr Pro Ala Pro Ala Lys Asp Ala Leu Arg Ala Leu
290 295 300

Thr Gln Arg Gly His Glu Ser Met Tyr Gln Gln Asn Lys Ser Val Ser

305	310	315	320
Thr Asp Glu Leu Gly 325	Pro Pro Gly Arg	Leu Glu Gly Val 330	Pro Arg Pro 335
Val Ala Arg Glu Lys 340	Thr Ser Leu Gly 345	Ser Leu Lys Arg	Ala Ser Val 350
Asp Val Asp Leu Leu Ala 355	Pro Arg Ser Pro Met 360	Gly Lys Glu Gly 365	Met
Val Thr Phe Ser Asn Thr 370	Leu Pro Arg Val 375	Ser Thr Pro Ser Leu 380	Asp
Asp Pro Ala Arg Arg 385	His Met Thr Ile 390	His Val Pro Leu Asp 395	Ala Ser 400
Arg Ser Arg Gln Leu Ile 405	Gly Glu Trp Lys 410	Gln Lys Ser Leu Glu 415	Gly
Arg Gly Leu Gly Leu Pro 420	Asp Glu Ala Ser Pro 425	Val His Leu Arg 430	Ala
Pro Ala Glu Gln Val Ala 435	Glu Glu Glu Glu Glu 440	Glu Glu Glu Glu 445	
Glu Glu Glu Glu Glu 450	Glu Glu Glu Glu Gly 455	Pro Val Pro Pro 460	Ser
Leu Tyr Pro Thr Val 465	Gln Ala Arg Pro Gly 470	Leu Gly Pro Arg Val 475	Ile 480
Leu Pro Pro Arg Pro 485	Gly Pro Gln Pro Leu 490	Val His Ile Pro 495	Glu Glu
Gly Val Gln Ala Gly Ala 500	Gly Leu Ser Pro Lys 505	Ser Ser Ser Ser 510	
Val Arg Ala Lys Trp Leu 515	Ser Val Ala Glu Lys 520	Gly Gly Gly Pro 525	Val
Ala Val Ala Pro Ser Gln 530	Pro Arg Val Ala Asn 535	Pro Pro Arg Leu 540	Leu
Gln Val Ile Ala Met Ser 545	Lys Ala Ala Gly 550	Gly Pro Lys Ala Glu 555	Thr 560

Ala Ser Ser Ser Ser Ala Ser Ser Asp Ser Ser Gln Tyr Arg Ser Pro
565 570 575

Ser Asp Arg Asp Ser Ala Ser Ile Val Thr Ile Asp Ala His Ala Pro
580 585 590

His His Pro Val Val His Leu Ser Ala Gly Ser Thr Pro Trp Glu Trp
595 600 605

Lys Ala Lys Val Val Glu Gly Glu Gly Ser Tyr Glu Leu Gly Asp Leu
610 615 620

Ala Arg Gly Phe Arg Ser Ser Cys Lys Gln Pro Gly Met Gly Pro Gly
625 630 635 640

Ser Pro Val Ser Asp Val Asp Gln Glu Glu Pro Arg Phe Gly Ala Val
645 650 655

Ala Thr Val Asn Leu Ala Thr Gly Glu Gly Leu Pro Pro Pro Gly Ala
660 665 670

Ser Glu Gly Ala Leu Gly Ala Gly Ser Arg Glu Ser Thr Leu Arg Arg
675 680 685

Gln Val Gly Gly Leu Ala Glu Arg Glu Val Glu Ala Glu Ala Glu Ser
690 695 700

Tyr Tyr Arg Arg Met Gln Ala Arg Arg Tyr Gln Asp
705 710 715

<210> 5
<211> 766
<212> PRT
<213> Rattus norvegicus

<400> 5

Met Gln Arg Ala Gly Ser Ser Gly Ala Arg Gly Glu Cys Asp Ile Ser
1 5 10 15

Gly Thr Gly Arg Leu Arg Leu Glu Gln Ala Ala Arg Leu Gly Gly Arg
20 25 30

Ala Val His Thr Ser Pro Thr Gly Gly Leu Gly Ala Arg Gln Val Ala
35 40 45

Gly Met Ser Ala Lys Glu Arg Pro Lys Gly Lys Val Ile Lys Asp Ser
50 55 60

Val Thr Leu Leu Pro Cys Phe Tyr Phe Val Glu Leu Pro Ile Leu Ala
 65 70 75 80

Ser Ser Val Val Ser Leu Tyr Phe Leu Glu Leu Thr Asp Val Phe Lys
 85 90 95

Pro Val His Ser Gly Phe Ser Cys Tyr Asp Arg Ser Leu Ser Met Pro
 100 105 110

Tyr Ile Glu Pro Thr Gln Glu Ala Ile Pro Phe Leu Met Leu Leu Ser
 115 120 125

Leu Ala Phe Ala Gly Pro Ala Ile Thr Ile Met Val Gly Glu Gly Ile
 130 135 140

Leu Tyr Cys Cys Leu Ser Lys Arg Arg Asn Gly Ala Gly Leu Glu Pro
 145 150 155 160

Asn Ile Asn Ala Gly Gly Cys Asn Phe Asn Ser Phe Leu Arg Arg Ala
 165 170 175

Val Arg Phe Val Gly Val His Val Val Gly Leu Cys Ser Thr Ala Leu
 180 185 190

Ile Thr Asp Ile Ile Gln Leu Ala Thr Gly Tyr Gln Ala Pro Tyr Phe
 195 200 205

Leu Thr Val Cys Lys Pro Met Tyr Thr Ser Leu Glu Gly Ser Cys Lys
 210 215 220

Glu Asn Ser Tyr Ile Val Glu Glu Ile Cys Ser Gly Ser Asp Leu Thr
 225 230 235 240

Val Ile Asn Asn Gly Lys Lys Ser Phe Pro Ser Gln His Ala Thr Leu
 245 250 255

Ala Ala Phe Ala Ala Val Tyr Val Ser Met Tyr Phe Asn Ser Thr Leu
 260 265 270

Thr Asp Ser Ser Lys Leu Leu Lys Pro Leu Leu Val Phe Thr Phe Ile
 275 280 285

Ile Cys Gly Ile Ile Cys Gly Leu Thr Arg Ile Thr Gln Tyr Lys Asn
 290 295 300

His Pro Val Asp Val Tyr Cys Gly Phe Leu Ile Gly Gly Gly Ile Ala
 305 310 315 320

Leu Tyr Leu Gly Leu Tyr Ala Val Gly Asn Phe Leu Pro Ser Glu Asp
 325 330 335

Ser Met Leu Gln His Arg Asp Ala Leu Arg Ser Leu Thr Asp Leu Asn
 340 345 350

Gln Asp Pro Ser Arg Val Leu Ser Ala Lys Asn Gly Ser Ser Gly Asp
 355 360 365

Gly Ile Ala His Thr Glu Gly Ile Leu Asn Arg Asn His Arg Asp Ala
 370 375 380

Ser Ser Leu Thr Asn Leu Lys Arg Ala Asn Ala Asp Val Glu Ile Ile
 385 390 395 400

Thr Pro Arg Ser Pro Met Gly Lys Glu Ser Met Val Thr Phe Ser Asn
 405 410 415

Thr Leu Pro Arg Ala Asn Thr Pro Ser Val Glu Asp Pro Val Arg Arg
 420 425 430

Asn Ala Ser Ile His Ala Ser Met Asp Ser Ala Arg Ser Lys Gln Leu
 435 440 445

Leu Thr Gln Trp Lys Ser Lys Asn Glu Ser Arg Lys Met Ser Leu Gln
 450 455 460

Val Met Asp Ser Glu Pro Glu Gly Gln Ser Pro Pro Arg Ser Ile Glu
 465 470 475 480

Met Arg Ser Ser Ser Glu Pro Ser Arg Val Gly Val Asn Gly Asp His
 485 490 495

His Val Pro Gly Asn Gln Tyr Leu Lys Ile Gln Pro Gly Thr Val Pro
 500 505 510

Gly Cys Asn Asn Ser Met Pro Ala Gly Pro Arg Val Ser Ile Gln Ser
 515 520 525

Arg Pro Gly Ser Ser Gln Leu Val His Ile Pro Glu Glu Thr Gln Glu
 530 535 540

Asn Ile Ser Thr Ser Pro Lys Ser Ser Ser Ala Arg Ala Lys Trp Leu
 545 550 555 560

Lys Ala Ala Glu Lys Thr Val Ala Cys Asn Arg Gly Asn Asn Gln Pro
 565 570 575

Arg Ile Met Gln Val Ile Ala Met Ser Lys Gln Gln Gly Val Leu Gln
580 585 590

Ser Ser Pro Lys Asn Ala Glu Gly Ser Thr Val Thr Cys Thr Gly Ser
595 600 605

Ile Arg Tyr Lys Thr Leu Thr Asp His Glu Pro Ser Gly Ile Val Arg
610 615 620

Val Glu Ala His Pro Glu Asn Asn Arg Pro Ile Ile Gln Ile Pro Ser
625 630 635 640

Ser Thr Glu Gly Glu Gly Ser Gly Ser Trp Lys Trp Lys Ala Pro Glu
645 650 655

Lys Ser Ser Leu Arg Gln Thr Tyr Glu Leu Asn Asp Leu Asn Arg Asp
660 665 670

Ser Glu Ser Cys Glu Ser Leu Lys Asp Ser Phe Gly Ser Gly Asp Arg
675 680 685

Lys Arg Lys His Ile Asp Ser Asn Glu His His His His Gly Ile Thr
690 695 700

Thr Ile Arg Val Thr Pro Val Glu Gly Ser Glu Ile Gly Ser Glu Thr
705 710 715 720

Leu Ser Val Ser Ser Ser Arg Asp Ser Thr Leu Arg Arg Lys Gly Asn
725 730 735

Ile Ile Leu Ile Pro Glu Arg Ser Asn Ser Pro Glu Asn Thr Arg Asn
740 745 750

Ile Phe Tyr Lys Gly Thr Ser Pro Thr Arg Pro Tyr Lys Asp
755 760 765

<210> 6

<211> 5083

<212> DNA

<213> Homo sapiens

<400> 6

ggatccacta gtaacggccg ccagtgtgct ggaattcgcc cttgaagcca ttgcagcaac 60

agcttgaggagg agggagctgg acgtcgtctc tcgccagaaa aacgggggagc aggagccaga 120

ctaaaggagg aagaggactg gcccgctcag ggaatagctg ggttgctgca aaaaggggagc 180

gggagaaggc gggggcgctg catgcagcgc gctggctcca gcggtggccg cggggaatgt 240

gacatcagcg	gcgccggg	cttggggctg	gaggaggcgg	ctgcctcag	ctgcgctgtg	300
cacacctcgc	ccgggggagg	acgcagaccc	gggcaggcgg	cagggatgtc	ggcgaaggag	360
aggccaaagg	gcaaagtgat	caaggacagc	gtcacccctc	tgccctgctt	ttatttcgtc	420
gagttgccta	tattggcatc	atcggtggtt	agcctctatt	tcctcgaact	cacagatgtc	480
ttcaaacctg	tgactcttgg	atttagctgc	tatgaccgga	gtcttagcat	gccgtacatt	540
gaaccaaccc	aggaggcaat	tccattcctc	atgttgctta	gcttggtttt	tgctggacct	600
gcaattacga	ttatggtagg	agaaggaatt	ctctactgtt	gcctctccaa	aagaagaaat	660
ggggctcggac	tagagcccaa	cattaatgct	ggaggctgca	acttcaattc	cttcctcaga	720
cgagctgtca	gattcgttgg	tgttcatgta	tttggtattat	gctctacagc	tctcattaca	780
gatatcatatc	agctgtccac	aggatatcaa	gcaccttact	ttctgactgt	gtgcaaacca	840
aactatacct	ctctgaatgt	atcttgcaaa	gaaaattcct	acattgtgga	agatatttgc	900
tcaggatctg	acctcacagt	tatcaacagt	ggcagaaagt	ccttccttcc	tcaacatgca	960
acccttgctg	cctttgcagc	tgtgtatgtt	tcgatgtact	tcaattccac	attaacggat	1020
tcctctaagc	ttctgaaacc	tctcttggtc	ttcacattta	tcctctgtgg	aataatctgc	1080
gggctaacac	ggataactca	gtataagaac	caccagttg	atgtctattg	tggcttttta	1140
ataggaggag	gaattgcact	gtacttgggc	ttgtatgctg	tggggaattt	cctgcccagt	1200
gatgagagta	tgtttcagca	cagagacgcc	ctcagggtctc	tgacagacct	caatcaagat	1260
cccaaccgac	ttttatctgc	taaaaatggt	agcagcagtg	atggaattgc	tcatacagaa	1320
ggcatcctca	accgaaacca	cagagatgct	agctctctga	caaatctcaa	aagagcaaatt	1380
gctgatgtgg	aatcattac	tccacggagc	cccatgggga	aggagaacat	ggttaccttc	1440
agcaatacct	tgccgcgagc	caatacccca	tctgtagaag	accctgtcag	aagaaatgcg	1500
agcattcatg	cctctatgga	ttccgctcga	tcaaagcagc	tcctcaccca	gtggaagaat	1560
aagaatgaaa	gtcgaaagtt	gtccttgcaa	gttatagagc	ctgagcctgg	gcagtcacca	1620
cccagatcca	tagaaatgag	gtcaagctca	gagccatcga	gggtaggggt	gaatggagac	1680
caccatggtc	ctggcaatca	gtacctcaaa	atccagcctg	gcgctgtccc	cggatgtaac	1740
aacagcatgc	ctggagggcc	aagagtgtcc	attcagtcctc	gtcctgggtc	ctcacagttg	1800
gtgcacatcc	ctgaggagac	tcaggaaaac	ataagcacct	cccccaaaag	cagctctgct	1860
cgggccaaagt	ggttaaaagc	tgctgaaaag	actgtggcct	gtaacagaag	caacagccag	1920
ccccgaatca	tgcaagtcat	agccatgtcc	aagcagcagg	gtgtcctcca	aagcagcccc	1980
aagaacactg	aaggcagcac	ggtctcctgc	actggctcca	tccgctataa	aaccttgaca	2040
gacatgagc	ccagtgggat	agtgagggtt	gaggctcacc	cagagaacaa	caggcccatc	2100

atacagatcc	cgtccactga	aggtgaaggc	agtggctcct	ggaagtggaa	agcccctgaa	2160
aagggcagcc	ttcgccaaac	ttacgagctc	aacgatctca	acagggactc	agaaagctgt	2220
gagtctctga	aagacagctt	tggttctgga	gatcgcaaga	gaagcaacat	tgatagcaat	2280
gagcatcacc	accacggaat	taccaccatc	cgcgtcaccc	cagtagaggg	cagcgaaatt	2340
ggctcagaga	cgtgtccat	ttctttcttc	cgcgactcca	ccctgcggag	aaagggcaat	2400
atcattctaa	tccttgaaag	aagcaacagc	cccgaaaaca	ctagaaatat	cttctacaaa	2460
ggaacctccc	ccacacgggc	ttataaggat	tgagtgatgt	ccattccatc	attagggcta	2520
ctcgcaaaag	accatatgtt	gattctacct	gtgttctgtt	ccagcgaatt	gggaagtctc	2580
accaagctag	attgtctacc	atcagcccag	aactctgtaa	cttttcagaa	ctgctatact	2640
caaacttgca	gatctcacat	caaggagagg	gaaaagcaca	atgcaagaac	ctaactaacg	2700
tgatgatatg	aagagttttc	ttaagacctg	tcgtcaaact	taaaagggtt	tgcagagggc	2760
agtatcaaaa	gaaagtgggt	ttcttcaa	gtatactatt	ttacttcctg	aatgtgccaa	2820
ctttggggat	ttttctttat	agtgagctgt	gggaaccag	aacacacacg	ttttccctac	2880
agcagaggcc	atgcagtatt	atatattcat	tttgagaat	ctgcacctac	agctcaatac	2940
gggtgggtgct	gattattata	gtacatatac	catgtaaact	ctcaaactct	atttagctgt	3000
gaaatagtgg	tgtgcaattc	cttgtaaag	aaatgctact	ttattaagaa	gatgctggct	3060
gctttgtgtt	agaataggac	accccgagc	ttctctgtag	tggctctgtc	acagtcaaaa	3120
aatgaaaagg	tttttgtg	tttcttcaa	attctgcttt	cttcaacatc	aaaaattgtg	3180
tagaaatatt	ttcagtga	gggaataact	agtacttttc	tgcatagttt	ttcttctgct	3240
tactttttat	ttaagtatag	gtactgctaa	tgaatctgtt	ttcttagtga	gtaaatttgc	3300
ataattttat	aaatattatt	ttagagaatc	ttttgaaatt	gttgtgatca	tattttgctt	3360
tctatggctt	ctccttaact	tattgattaa	ttttttgaag	ttatagatat	gttctcctat	3420
tttaaaagca	aaaataacaa	ttgacattcc	ttgagcaaaa	tatactgctg	tgaatttgca	3480
aacaagaaat	ctgagccaaa	acttgacatt	gtgggttaca	ttgccagaaa	tgttggtcaa	3540
gtttgccctt	agatgtctac	aactagctgg	cataggttgc	catcttaaca	agtaatctaa	3600
aagtcccat	cggttctaca	ttattaactt	tttttttcta	tatcctgatg	accagtaaat	3660
tagagccaca	ctggttaagt	ttgactcgtc	tctaaaacgt	ttttgttaat	tggacaccaa	3720
gaggaagaat	ctgaaaaaaaa	aatgcatgtt	ggtaagtaaa	agtatctcac	ggtacaaatt	3780
aagaatgact	ttcttcaaaa	tatctgaata	ggtgcagttt	tagtttaaca	tgcaaacac	3840
cattgttgct	acctatcctg	aatcaagcct	tgagcctaaa	tcaaagcaaa	ccaataccat	3900
tgataagaag	aagataaaaa	caaaatattt	tggagtgttt	tccaacttaa	agtatgaaga	3960
catactcagt	tcttggaact	tagtattaaa	ccttttttat	gccatttcat	aagaattccg	4020

atatatactt gatgattgcc aaggggatga aaggaaacaa cagagatggt tgatctgatc	4080
ttagctcact ttccaataac agaaggagtt gtttacagat gaatagtatc acatcattat	4140
caatttccac atgaaaaagg tggagctttc tagaaaaacc aacctctaag gcattaggaa	4200
tttagctgaa accagcagaa ttgaaaactc tggcaataaa acatggactc aaccatatcc	4260
cttctggcaa tttccttctc agagagggga gtgggaataa aatgttgctt tccccacttc	4320
tcaccaccac cgccatcatg acgctcatac tggcttttgc ctgtttgtag aggaaaagggt	4380
gggctggttt tagtactctg aaggacaaaa acaagcaaac aaaaaccctt gctgcagcat	4440
ttcaggtgca gtatgatatt tcctaattctt tcctatttct taacaaaaga ttttaaagta	4500
cttctctagt cattgaagtt tttttttctt tacataaata ttgatataatt ctttttctac	4560
tcaaagtgcc aaaggctaca gtttttaatg acttaacaaa ttgtaccaca ttgttaagga	4620
catataatga tagacactag aactcagacc tctgcatgta tatttgataa catgtctttt	4680
gtaaaacaaa aattacaaaa aaatttgttt acattccact ggtaccttaa tttaaaataa	4740
atcagactaa aagggtggtat ctcttcttag tgttctatct atcttatttg ctaatgggag	4800
cacttcttcc tttgttaggc tgtgctttac tgataaaacc aagtattgaa taaagagagt	4860
taattatctt tttaaagtaa ataaaattat gaaaatatat atagtatata taaagtcttg	4920
tgtttaaaaa aatgttatgc aatgttttcc aaactgataa agtttgtaaa gtgctataaa	4980
tgtattttgt taagtacaga taaaagctat tgtgtgagta tattgtgcta aaatcataga	5040
aataaagatt agatttcttc atcaaaaaaa aaaaaaaaaa aaa	5083

<210> 7

<211> 2356

<212> DNA

<213> Homo sapiens

<400> 7

aggagacgcg ctttgtgctg ggcgcgggcc gcgccagcca cggcctgcgg cgcccgcggc	60
accatgatct ccaccaagga gaagaacaag atcccgaagg acagcatgac gcttctgccc	120
tgcttctact tcgtggagct gcccatagtg gcttcttcca tcgtatcctt gtacttcttg	180
gagctgaccg acctcttcaa gccggccaag gtgggcttcc agtgctatga ccgcactctc	240
tccatgccct acgtggagac caacgaggag ctcatcccg cgtctgatgct gctcagcttg	300
gccttcgcgg cccctgccgc ctcgatcatg gtggccgagg gcatgttgta ctgtctgcag	360
tcccggtgtg ggggccgtgc cggggggccc gccggggcgg agggcagcat caacgccggc	420
ggctgcaact tcaactcctt cctgcggcgt acgggtgcgg ttgtgggtgt ccacgtgttc	480
ggcctgtgtg ccacagccct ggtgacggac gtgatccagc tggccacggg ttaccacact	540
cccttcttcc tcaccgtctg caagcccaac tacactctcc tgggcacgtc ctgcgaggtc	600

aaccctaca tcacgcagga catctgctcc ggccacgaca tccacgccat cctgtctgca	660
cggaagacct tcccgctcca gcacgccacg ctgtcagcct tcgccgcggt ctatgtgtcg	720
gtgagtcggg cacctcactg cccttcccag gccctcttgc tgacctgtgg ggagccctcc	780
ctgaccccaa ccccatgcc ccagatgtac ttcaactcgg tcatctcgga caccaccaag	840
ctgctgaagc ccctcctggg ctctgccttt gccatgcgcg cgggcttatg cgggctcacg	900
cagatcacgc agtaccgcag ccaccctgtg gacgtgtatg ccggcttctt catcggggcg	960
ggcatcgctg cctacctggc ctgccacgcg gtgggcaact tccaggcccc acctgcagag	1020
aagcccgcgg ccccgggccc cgccaaggac gcgctgcggg ccctgacgca gcggggccac	1080
gactcggttt atcagcagaa taagtcggtg agcacgcgag agctggggcc ccagggcg	1140
ctggagggcg cgccccggcc cgtggcccg gcagaagacct cgctgggcag cctgaagcgc	1200
gccagcgtgg acgtggacct gctggccccg cgcagcccca tggccaagga gaacatggtg	1260
accttcagcc acacgctgcc cagggccagc gcgcctcgc tggacgacct cgcgcgccgc	1320
cacatgacca tccacgtgcc gctggacgcc tcgcgctcca agcagctcat cagcgagtgg	1380
aagcagaaga gcctggaggg ccgcggcctg gggctgcccg acgacgccag ccccgggcac	1440
ctgcgcgcgc ccgccgaacc catggcgag gagggagga agggaggagga cgaaggagaa	1500
gaggaggagg aggaagagga ggaggacgag ggcccggccc cgccctcgt ctacccacc	1560
gtgcaggcgc ggccggggct ggggcctcgg gtcctctcc caccgcgcgc ggggcccgcg	1620
ccgctggtgc acatcccga ggaggcgcg cagacggggg ccggcctgtc cccaaaagc	1680
ggcgccgggg tgccgcgcaa gtggctcatg atggccgaga agagcggggc ggcagtggcc	1740
aaccctcgc ggctgctgca ggtcatgcc atgtccaagg ctccgggcgc gccgggcccc	1800
aaggcgggcg agacggcgtc gtcgtccagc gccagctccg actcctcgca gtaccggtcg	1860
ccgtcggacc gcgaactcgc cagcatcgtg accatcgacg cgcacgcgc gcaccacccc	1920
gtggtgcacc tgtcggccgg cggcgcgcc tgggagtga aggcggcggg cggcggggcc	1980
aaggcgagg cgacggcg ctacgagctg ggggacctg cgcgcggtt ccgcggcggg	2040
gccaagcccc cgggctgtc ccccggtcg tcggtcagcg acgtggacca ggaggagccg	2100
cggttcgggg ccgtggccac cgtcaacctg gccacgggcg aggggctgcc cccgctgggc	2160
gcggccgatg gggcgctggg cccgggcagc cgggagtcca cgctgcggcg ccacgcgggc	2220
ggcctggggc tggcgagcg cgaggcgag gcggaggccg agggctactt ccgcaagatg	2280
caggcgccc gcttccccga ctacgcggc ggggcccggg gcgggcccgg ggcgggccga	2340
gggcgcgggc ggccgc	2356

<211> 5708
 <212> DNA
 <213> Mus musculus

<220>
 <221> misc_feature
 <222> (460)..(460)
 <223> n is a, c, g, or t

<400> 8
 ygagc gatga tgccccat tt accctttctc ttcagatgca ggaaat tttc actctgttcc 60
 ccagctgatt ggagcttttt ctaggtgctt ccctgggagt tacctcccta gagatcagca 120
 ggcagggctg tcacgcttgg gtagcagcca gctcccagtg aattccttct gtggcctact 180
 tgtccttatg aagtccgagt tttaattttg cacaggtagg aggtctcttt tgctatggat 240
 agggcgata acggtgctac cattagaaaa caggcttctg ttttctagga aggcaagagg 300
 aaccccaggt aggggacctt gtgagaccag gtgacttggc tcctcagcct tgcttctaca 360
 gaaaccagga gtgcttcccc ccactcttcc ctatttttga cgtcaagctc aaccagccag 420
 cagaggagcc tcacggcttg ggcggtggag agagagcccn aggagagtgg caggggagggg 480
 aagccatctc agcaacagct tggagaggga gctgctatcc cttgcccgca aaacacggac 540
 taaagccagg ctgaagaaga cctgcgggct cgggctcggg gatccgcggg gttactgcaa 600
 agaagggcgg ggaaaaggcg ggggcgctgc atgcaacgcg ctgggtccag cggtgcccgc 660
 ggggaatgtg acatcagcgg cgccggggcg ttgcggctgg agcaggcagc tcgcctcggg 720
 ggccgcacgg tgcacacctc gcccggggga ggacttggag cccggcaggc ggccgggatg 780
 tcggcgaagg agaggccaaa gggcaaagt atcaaggaca gcgtcacctt cctgccctgt 840
 ttttatttcg ttgagttgcc tatattggca tcatcagtg ttagcctcta cttcttggaa 900
 ctcacagatg tcttcaaacc tgtgcactct ggattcagtt gctatgatag gagtcttagc 960
 atgccgtaca ttgagccaac ccaggaggcc ataccattcc ttatgttgct tagcttggct 1020
 tttgctggac ctgcaattac gatcatggtg ggtgaaggga ttctatactg ctgcctctcc 1080
 aaaagaagaa acggagctgg attggagcct aacatcaacg ccggaggctg caacttcaac 1140
 tcctttctca ggagagccgt cagattcggt ggtgtccatg tgtttggact gtgctccaca 1200
 gctctcatta cagatatcat acagctctcc acaggatatc aggcaccata ctttctgact 1260
 gtgtgcaagc caaactatac ctctctgaat gtatcctgca aagaaaactc ctacatcgtg 1320
 gaagatattt gttcaggatc tgaccttaca gtcacaaaca gtggcagaaa gtcattccca 1380
 tccaacatg cgaccctcgc tgcctttgcc gctgtgtatg tgtccatgta cttcaattcc 1440
 acattaaccg attcctctaa gctcctgaaa cctctcttgg tcttcacatt tatcatctgt 1500
 gggatcatct gcggactaac acggataact caatataaga accatccagt cgatgtctat 1560

tgtggctttt taataggagg aggaatcgca ctatatattgg gcctgtatgc tgtagggaat	1620
tttttgccta gtgaagacag tatgcttcag cacagagatg ccctcaggtc actgacagac	1680
ctcaatcaag accccagcag ggttttatca gctaaaaatg gtagcagtgg tgatggaatt	1740
gctcacacag agggatcct caaccgaaac cacagggatg caagctcctt gacaaatctc	1800
aagagggccca acgctgacgt agaaatcatc actcctagga gcccctatggg gaaggaaagc	1860
atggtgacct tcagcaacac gctgcccagg gccaacaccc cctccgtgga agaccagtg	1920
agaagaaatg cgagcatcca tgcctctatg gattctgccc ggtccaaaca gctccttacc	1980
cagtggaaga gcaagaatga gagtcgtaag atgtccctac aggttatgga cactgaacca	2040
gaaggccagt caccaccag gtccatagaa atgaggcca gctcagagcc ctcgagggtg	2100
ggggtgaacg gagatcacca tgtccctggc aatcagtacc tcaagataca gcctggcaca	2160
gtccccgggt gcaacaatag tatgccggga gggccacgcg tgtccatcca gtcccgccct	2220
ggctcttccc aattggtgca catccccgag gagaccaggg aaacataag cacctcgccc	2280
aagagcagtt ctgcgcgagc caagtggctg aaagcagctg agaagaccgt ggactgtaac	2340
cggagcaaca accagccacg catcatgcag gtcacgcca tgtccaagca gcagggcgtg	2400
ctgcagagca gcccgaagaa tgccgaaggt agcactgtca cctgcacagg ttccatccgc	2460
tacaaaaccc tgactgacca tgagcccagc ggcacgtgc gagtggaggc tcatcccgag	2520
aacaacaggc ccatcattca gatcccgctg tccactgagg gtgaaggcag cggctcctgg	2580
aagtggaaag ttccggagaa aagtagtctg cgccaaacct atgagctcaa cgacctcaac	2640
agggactcag aaagctgtga gtccctcaaa gacagctttg gttctggaga tcgcaaaaga	2700
agcaacatcg acagcaatga gcaccaccac cacggcatca ccaccatccg agtgacccccg	2760
gtggagggca gcgagatagg ctgagagaag ctgtccgtgt cctcctcacg cgactccacc	2820
ctgcgcagga agggcaacat catcttgatc ccggaaagaa gcaacagccc tgaaaacaca	2880
agaaacatct tctacaaagg aacctcccc acgcgggctt ataaggattg agagatggcg	2940
gcccttcttg tcatcatttt gatgacaccc ccacctcccc atccccacc ctcaccccaa	3000
gaccactcgt ttattgtacc ttgtgctctt ttgggttttt tgttttgttt tgtttggggg	3060
cctttttttt tccctagaag atatggagag ccttcttgct caactagatt gttcaccatc	3120
agcctggaac tctcactgaa ccaccacaga aatcgtggcg attttacacc aagggaaggg	3180
aaaagcacia agcaagaccc gaactaaact catcatcaga acagttctta agacacaggc	3240
tttgagaag gtagtattaa gataaagtgg tttcctccga tgtatagtat ttaactttct	3300
gaatgtgcca acttaatgga gttttttttt ttcattataa ttagctgtgg gaacccaaaa	3360
cacataggtt ttcccaacag cagaggccat gcggtattat atattattca tttttgcaga	3420
ctctgcacca gaagagcaga ctgggtggtg ctgattatca cagtgcattc accatttaaa	3480

ctctcaaact ctatgtagct gtgaaatagt ggtgtgcaac tcctcgtcag agaaatgcta	3540
cttcattcag aagacgccag tgactttgtg ttagaataga ccattcttgg cttccctgta	3600
gtggctctct cacagttgaa aagaaaagaa aagaaaaaga aaaagaaaga gagagagaga	3660
aagagagaaa gaaagaaaga aagaaagaaa gaaagaaaga aagaaagaaa gaaagaaaga	3720
aagaaagaaa gaaagaaaga aagaaagaat tggatgaatt ggacagggct ttgagcattt	3780
ctttgaaaga tgcttttttt caacatctga aagcttgtag gaatgttttc agtgaaacag	3840
aataactagt tctctgcac gtttttcttc tttttattta agtattggta atgctgcttt	3900
ctgggtttttt gttttttgtt ttagtgagtg catttgcata tttaaaatac attgttttag	3960
agaatatttt gaaattatta tgattacatt ttccatttta tggctttacc ttagtttatt	4020
aagttttctg agggttacac atattcttct attttaagaa agcaaaaagt acaacttgca	4080
ttctttgtgc aaaatacact gctgtgaggt cctacactag aaatctgagc caaaggttga	4140
aactgtgcgt gccaatgcca gatacgctgg tcaagggtcaa gatgtctcca atccgatggc	4200
ataggttatc acatcagtaa gtaatccaa aatttcattt tgttccagag catttcattt	4260
tcatgttatc ttgataatca ccatattgga gccacagtgg gggtgagttt gactcccttt	4320
cctgacacac ttttaactgc acaccaacag taagaatcta ggcaaagtct aattgataaa	4380
tagatgtgta tcacagtata agtttagaaa gcatatcttc aaaatgtcag accaggtaaa	4440
gctttcgtgc ttagagtata accaacagtt ttggatgtct gtcttgaatc tagaacctta	4500
agcctaaatc aaaggaaacc ttactgttga tagcaagaag ataacaacat atttttgaag	4560
tggttttcca agctagctgt ttaaagtgtg gagaaggatt tggttcttga aatttggtat	4620
taaccttttt tcatgccatg tcttaagagt tataatgtac actcgatgat tgccaagaga	4680
gggggagggg gaggaaaaca gccaacagca gagctgggtg gtctgaactc agtgcagttt	4740
tcaatgagaa caacagctgt ccagcaagga atcatatcat ccattctcag cttctacatt	4800
caaagggcag agcttttttag aaaactcaac ctctaaggc attaggaatt gagctgaaac	4860
cagcagaatt gaaaactctg gcaataaaat atagactcaa tcgtaaccct tctggcaagt	4920
tccttctcag agaaggaagt gggagtaaaa tgtggccttc cccacttctt tacatcaccc	4980
ctgtcacaat gtccccgctg gcctggccag tttcgagagg gaaggggtgga ctggtttttag	5040
tactctgaag aaaacccaag ctgcagtatt tgaggtgcag tataatattt cctaattctt	5100
cctatttctt aacaaaaaaa gattttaag tacttctcta ctattgaat ttttgttctt	5160
tacatactat tgatatattc tttttctact caaaagtgcc aaaggctaca gtttttaatg	5220
acttaacaaa ttgtaccaca ttgttaagga aatataatga tagacactag aattcagacc	5280
tctgcatgta tatttgataa cacatctttt gtaaaaaata aataattaca aaaaatttgt	5340

ttacattcca caggtacctt aattttaa	aat	aatcagact aacaggtggt atctcttctt	5400
agtgttctat ttatcttatt tgcta	atgag aacaattctt cttctgttag	gctgtgcttt	5460
attgataaaa ccaagtattg aataa	aggga gttaattatc tttttaaa	agt aaatgaaatt	5520
ataaatatat aatatatata agt	attgtg ttttaataaaa tgttatg	caa tgttttccaa	5580
actgataaag tttgtaaagt gct	tataaatg tattttgtta agtac	agatc aaagctatcg	5640
tgtgagtata ttgtgctaac at	catagaaa taaagattag attt	cttcat caaaaaaaaa	5700
aaaaaaaa			5708

<210> 9
 <211> 2783
 <212> DNA
 <213> Mus musculus

<400> 9	
ggcgcggggc gcgggatctc gccttcccca cactgcgccg	cgcgctccta ggtccccgag 60
atcaaccgtg ctgcgcgaag gcgcacagaa ggcacgcgca	aggggcaggg acgagacca 120
gattgcggtc cccgcgcagg gccagagacc ctgagacc	cg cagagtggga ggcggaggcg 180
cccaagccgg gggcgggggtt ggggcggggc cgagcggg	ct gcagagcgcg cagacaaaga 240
gctcacgcgc cgcacaccgc gcccgggcct cgcgtcc	ctc ccacgcgggg accctggctt 300
tgtgtccgcg ctttgcaagg gactccgcc tcttgtct	cc ctccagggtga acgagccgcg 360
tgcgcgcgcg ggggctccgg cctctccctt cccaagt	tcc cccctctgct ccagcctcag 420
tttccccgag ccctcttggc taacccttat ctcccc	acct ggagactcca tctgcgcagc 480
tcctagtctc cagcctgcct ttgtccccct gcaatct	cgg cctacagcgc caacgtcacc 540
atgcttgcta tgaaggagaa gaataaaacg ccgaagg	aca gcatgacact cctgccttgc 600
ttttatttcg tggagctacc catcgtggt tcttcc	attg tgtctctgta cttcctggag 660
ctgactgacc tgttcaagcc ggccaagggt ggcttt	cagt gctacgaccg cgccctgtcc 720
atgccctatg tggagacaaa tgaagagctg atccc	attgc tcatgctgct gagcctggcc 780
ttcgtctcac ctgcggcctc aatcatggtc ggcgag	ggca tgggtctactg tctgcagtcc 840
aggctctggg gccgtggtcc agggggcggtg gaggg	cagca tcaacgctgg tggctgcaac 900
ttcaactcct tcttccggcg cacagtgcgc tttgt	gggtg tacaagtgtt tggcctgtgt 960
gccacggctc tggtgacaga tgtcatccag ctggcc	accg gctaccacac acctttcttc 1020
ttaacgggtc gcaaacccaa ttacaccctg ctgggc	actt cctgtgagtc gaacccttac 1080
atcacacagg acatctgctc tggccacgat acccat	gcca tctgttcagc aaggaagacc 1140
ttcccgtccc agcacgccac tctgtcagcc tttgt	gctg tctacgtgtc gatgtacttc 1200
aacgcgggta tctcggacac cacgaagctg ctga	agccca tcttgtgtt tgccttcgcc 1260


```

attgctgcgg gcgctctgcg cctcacacag atcaccagct atcgaagcca ccctgtggac 1320
gtctacgctg gctttcttat cgggtgctggc atcgctgcct acctggcctg ccacgctgtg 1380
ggcaacttcc aggcaccacc tgcagaaaag gttcctaccc cagctcctgc caaggacgcc 1440
ctgcgagcgc tgacacagcg gggccatgag tccatgtatc agcagaataa gtctgtcagt 1500
accgatgagc tggggccctcc agggaggcta gagggcgtgc ctggcctgtg ggctcgagag 1560
aagacatctc ttggcagctt gaagcgagcc agcgtggatg tggacctgct ggccccacgt 1620
agcccatgag gcaaagaagg catggtcacc ttcagcaaca cactgccccg ggtcagcacg 1680
ccctcgctgg atgacctgc acggcgccac atgactatcc acgtgcccct tgatgcctcc 1740
cgttccaggc agcttatcgg tgagtggagc caaaaatccc tggagggacg tggcctgggt 1800
ctgcctgatg aagccagccc tgtgcatctg agggccccag cagagcaggt agcagaggag 1860
gaagaggaag aggaggagga ggaggaagaa gaggaggaag aggaggaaga ggaagggcct 1920
gttcaccct cactctaccc tactgtccag gctcgccag gccttgggcc ccgagtcac 1980
ctgcctccaa ggccagggcc ccagcccctc gtacacatcc ctgaggaagg cgttcaggct 2040
ggagctggcc tgtcacccaa gagcagcagc tcatcagtga gggccaagtg gctgtcagt 2100
gctgagaagg gtggggggccc agtggtgtg gctccatccc agccccgggt ggccaacca 2160
cccaggctac tacaggctcat cgctatgtcc aaggcggcag ggggccccaa ggctgagaca 2220
gcttcgtcct ccagtgccag ctccgactct tcccagtaca ggtccccgtc agaccgtgac 2280
tctgccagca ttgttacaat cgatgcccac gcacctcacc atccagtgggt gcacctgtct 2340
gctggcagca caccctggga gtggaaggct aaagtagtgg agggtgaggg tagctatgag 2400
ctgggggacc tggcacgcgg cttcagaagc agctgtaaac agcctggaat gggcccgggg 2460
tctccagtca gtgatgtgga ccaggaagaa ccccggtttg gggcagtggc cactgtcaac 2520
ctggccactg gggagggctt gccccacca ggtgcaagtg aaggggccct aggtgcaggc 2580
agcagagaat ccaccctaag gcgccagggt ggcgggctgg cagagagaga agtggaggct 2640
gaggcagaaa gttattatag gaggatgcag gccaggaggt accaggacta agcctggcaa 2700
aactgtgatg gggattaggg gagctggggg tgtctacagc cccaccaata aaaacttatg 2760
ctattaataaa aaaaaaaaaa aaa 2783

```

```

<210> 10
<211> 5372
<212> DNA
<213> Rattus norvegicus

```

```

<220>
<221> misc_feature
<222> (1137)..(1137)
<223> n is a, c, g, or t

```

<220>
 <221> misc_feature
 <222> (2899)..(2899)
 <223> n is a, c, g, or t

<220>
 <221> misc_feature
 <222> (2917)..(2919)
 <223> n is a, c, g, or t

<220>
 <221> misc_feature
 <222> (2921)..(2924)
 <223> n is a, c, g, or t

<220>
 <221> misc_feature
 <222> (2929)..(2929)
 <223> n is a, c, g, or t

<400> 10
 atgccccaat tcacctctc tcttcagatg caggcaaaaa attctcactt tgttcctcgg 60
 cggattggag ctttttctag gtgcttcct tggagttacc tccctagaga tcagaaggtc 120
 agggctgtca cgcttgggta gcggccagct ccagtgaaat tccttctatg gcctacttgt 180
 cagtatgaaa tctgagtttt aattttgcac aggtggaggt ctcttttgct atgggtaagg 240
 tggatagcgg tgcccagcaa gtcctgctc tctagaaagg caacaagaac ccaggtagg 300
 agaccaggtg ccttggtcc tcagccttg tctgcagaa accaggagt cctcccccca 360
 ctatttttga cgtcaagctc agacaaccag cagaggagcc tcacagcttg ggcggtggag 420
 agcccaggga gagtggcagg gaggggaagc catcacagca acagcttgga gggggagctg 480
 gctatcactt tccctcaaaa tacggactaa aaccgggtg aagaagacct gcgggatcag 540
 ggaagcgccg ggttactgca aagaagggcg gggaaaagga gggggcgctg catgcagcgc 600
 gctggctcca gcggtgcccg cggggaatgt gacatcagcg gcaccgggcg cttgcggctg 660
 gagcaggcag ctgcctcgg tggccgcgca gtgcacacct caccacggg aggactcgga 720
 gccgggcagg tggccgggat gtcggcgaag gagaggcaa agggcaaagt gatcaaggac 780
 agcgtcaccc tctgccttg tttctatttc gttgagttgc ctatattggc gtcacggtg 840
 gttagtctct acttcctgga actcacagat gtcttcaaac ctgtgcactc cggattcagt 900
 tgctatgaca ggagtcttag catgccgtac attgagccaa cccaagaggc cattccattc 960
 ctcatgttgc ttagcttggc ttttgctgga cctgcaatta cgatcatggt ggggtgaaggg 1020
 attctgtact gctgcctctc taaaagaaga aatggagctg gattggagcc caacatcaac 1080
 gccgggggct gcaacttcaa ttcctttctc agaagagccg tcagatttgt tgggtgncac 1140
 gtggttggaac tgtgtccac agctctcatt acggatatca tacagcttgc cacaggatat 1200
 caggcaccat actttctgac tgtgtgcaag ccaatgtaca cctccctgga aggatcctgc 1260

aaggaaaatt cctacatcgt ggaagaaatt tgttcgggat ctgacctaac agtcatcaac	1320
aatggcaaaa agtcattccc gtcccagcat ggcaccctcg ctgcctttgc tgctgtgtat	1380
gtgtccatgt acttcaattc cacattaact gactcctcca agctcctgaa acctctcttg	1440
gtcttcacat ttatcatctg tggaatcatc tgcgggctaa cacggatcac tcagtataag	1500
aaccatcccg tcgatgttta ttgtggcttt ttaattggag gaggaatcgc gctgtatttg	1560
ggcctgtatg ctgtagggaa ttttctgcct agtgaagaca gcatgcttca gcacagagat	1620
gccctcaggt cactgacaga cctcaatcaa gaccccagca gggttttgtc agctaaaaat	1680
ggtagcagcg gtgatggaat tgctcacaca gagggtatctc tcaaccgaaa tcacagggat	1740
gcaagctcct taacaaatct gaagagggcc aacgctgacg tagagatcat cactcctagg	1800
agccccatgg ggaaggagag catggtgacc ttcagtaaca cgctgcccag ggccaacacc	1860
ccgtccgtgg aagaccgggt gagaagaaac gcgagcattc acgcctctat ggattctgcc	1920
cgggtccaagc agctcettac ccagtggaag agcaagaacg agagtcgcaa gatgtccctg	1980
caggttatgg actctgaacc agaagggcag tcaccaccca ggtccataga aatgaggtcc	2040
agctcagagc cctcgaggggt ggggggtgaat ggagatcacc atgtcccggg caatcagtac	2100
ctcaagatcc agcctggcac agttcccggg tgcaacaata gtatgcctgc agggccacgt	2160
gtgtccatcc agtcccgccc tggtctcttc caattgggtgc acatccccga ggagaccag	2220
gaaaacataa gcacctcgcc caagagcagt tccgcacggg ccaagtggct gaaagcagca	2280
gagaagaccg tggcctgtaa ccggggcaac aaccagccac gcatcatgca ggtcatcgcc	2340
atgtccaagc agcagggcgt gctgcagagc agccccaga atgccgaagg tagcacggtc	2400
acctgcacag gctccatccg ctacaaaacc ctgactgacc acgagccgag tggcattgtg	2460
cgagtggagg ctcatccga gaacaacagg cccatcatcc agattccatc gtccaccgag	2520
ggcgaaggca gtggctcctg gaagtggaaa gctccggaga agagtagcct gcgccaacc	2580
tatgagctca acgacctcaa cagggactcg gaaagctgcg agtccctcaa agacagcttt	2640
ggttctggag atcgcaagag aaagcacatc gacagcaatg agcaccacca ccacggcatc	2700
acaaccatcc gcgtgacccc agtggagggc agcgagatag gctcagagac gctgtctgta	2760
tcctctcac gcgactccac cctgcgcagg aagggaaca ttatcttgat ccctgaaaga	2820
agcaacagcc ctgaaaacac aagaaacatc ttctacaaag gaacctcccc cacgaggcct	2880
tataaggatt gaatccccnc ccttctgcct catcatnnng nnnncatcnc ccaccccccc	2940
acccagcca cacatttatt atagtacctt gtgctctttt gggatatttg ttgttgtttt	3000
ggttttggtt ttttgttttc ttttctagtg gatattggaga gccttcttgc ccaactagat	3060
tgttcaccat cagcctggaa ctctcactga cccaccacag aaaccgtggg gatthttccac	3120

caagggaaag gaaaagcaca aagcaagacc cgaactaatc tcatcatctg aatagttctt	3180
aaggcctgtc actgagactt cgcagagggc agtattagga taaagtggtc tcctccgatg	3240
tatagtatth aactttctga atgtgccaac tttgaagttt cttttttttt tttttcatta	3300
taattagcgg tgggaaccca aaacactaca ggtttccccg gcagcagagg ccacgcagta	3360
ttatatagta tttatthttg cagactctgc accagaagag cagactgggc ggtgctgatt	3420
accacagtac atctaccatt tcaactctca gactctgtgt agctgtgaga gagtgggtgta	3480
caacccccctt gtcagagaaa tgctacttca ttcagaagac gccagtgact ttgtggtagc	3540
ataggccatt cttggcttcc ctgtagtggc tctctcacag tcaaaaagaa aaaagaaaga	3600
aaaaactgga caggggtctg agcatttctt tgaaagattc tttgttttca acatttgaaa	3660
gtctgtaggg atgttttcgg tgagacggaa caactagttc tccgcacgt ttttctgctt	3720
tttatttaag tattggtaat gctggtttct ggtatctttt ttttttagtg agtgcatttg	3780
catatttaaa atacattggt ttagagaata ttttgaaatt attataatca cattttccat	3840
tttggtgctt catcttagtt tattaagttt tttgaggtta tgcacaccat tcttctatth	3900
taagaaagca acagtgacca tttacattcc ttgtgcaaaa tacactgctt cgaggtccta	3960
cgcaagaaat atgagccaac ggttgaaact gtgcgttcca ttgccagata cattgggtcac	4020
attcaagatg tctccaatcc gatggcatag gttatcacat cagtaagcga tccccaaact	4080
tcattttctt ccagagcatt tcatttttgt gttatcctga taatcaccat atcggagcca	4140
ccatgggggt gagcttgact cccttccctg acacactttt aactggacac caacagtaag	4200
gacctaggca aaggctaatt gatagataga agtgtatcac agcatacatt tagaaagtgt	4260
atcttcaaaa cagcagacca ggcaaagctt ttgtgcttag agtactattg acagttttgg	4320
ctgcctgtct tgaatctaga accttgagcc aaaatcaaag gaaaccatac tgttgatagc	4380
aagaagatta ccacaagata tttttgaagt gttttcccag gttagatggt taaagagtgg	4440
agacatatat ttggtccttg aaatttgta taaaacctt ctcatgccat gtcttaagaa	4500
ttataatgga cacttgatga ttgccaagag ggggagcaaa aacagcgaac agcagagctg	4560
gttggtctga actcagtga gctttcagt agaacaaaag ctgtccagca aagagtcaca	4620
tcattcattc tctagcctct ccattcaaag ggcagagctt tttttttttt ttaagaaaac	4680
tcaacctctt aaggcattag gaatttagct gagaccagca gaattgaaaa ctctggcagt	4740
aaaatatagg ctcaaccata accttcttg aaagttcctt tttttttttt ttttctttt	4800
ttttattttt tttttttttt ttgtgatcga gttgactgca gtagttcatc atattctttt	4860
tctactcaaa gtgccaaagg ctacagtttt taaacgactt aacaaattgt accacactgt	4920
taaggaacta taacgataga cactagaatt cagacctctg catgtatatt tgataacaca	4980
tcttttgtaa aaaataaata attacaaaaa atttgtttac attccactgg taccttaatt	5040

taaaagaaat cagactaaca ggtggtatct cttcttagtg ttctatttat cttacttgct 5100
 aacgagagca attcttcttt tgtaggctg tgctttattg atgaaaccga gtattgaata 5160
 aagagagtta attatctttt taaagtaaata aaaactatga atatataata tatatgaagt 5220
 atcgtgttta ataaaaatgt tatgcaatgt tttccaaact gataaagttt gtaaagtgc 5280
 ataaatgtat tttgttaagt acagatcaaa gctatcgtgt gagtatattg tgctggcatc 5340
 atagaaataa agattagatt tcttcatcaa aa 5372

<210> 11
 <211> 30
 <212> DNA
 <213> Rattus norvegicus

<400> 11
 caccacagct gagagggaaa tcgtgcgtga 30

<210> 12
 <211> 30
 <212> DNA
 <213> Rattus norvegicus

<400> 12
 atttgcggtg cagcatggag gggccggact 30

<210> 13
 <211> 51
 <212> DNA
 <213> Rattus norvegicus

<400> 13
 gcagaggtct gaattctagt gtctatcggt atagttcctt aacagtgtgg g 51

<210> 14
 <211> 14
 <212> PRT
 <213> Artificial

<220>
 <223> Peptide representing amino acids 490 to 507 of Rattus norvegicus
 PRG-1 protein with an added amino terminal cysteine

<400> 14

Cys Val Gly Val Asn Gly Asp His His Val Pro Gly Asn Gln
 1 5 10

<210> 15
 <211> 26
 <212> DNA
 <213> Rattus norvegicus

<400> 15
 ctaggcttgt agctgtgggg aatttc 26

<210> 16
<211> 22
<212> DNA
<213> Rattus norvegicus

<400> 16
tcaatcctta taagcccgtg tg 22

<210> 17
<211> 20
<212> DNA
<213> Rattus norvegicus

<400> 17
gaactttgcg agtgagctgg 20

<210> 18
<211> 20
<212> DNA
<213> Rattus norvegicus

<400> 18
tgcggagagc tttaacctcc 20

<210> 19
<211> 20
<212> DNA
<213> Rattus norvegicus

<400> 19
cctacctctt cctcatgttc 20

<210> 20
<211> 20
<212> DNA
<213> Rattus norvegicus

<400> 20
taaagggtgg agtccatcag 20

<210> 21
<211> 21
<212> DNA
<213> Rattus norvegicus

<400> 21
ggaattgcct ctgcaacatc t 21

<210> 22
<211> 18
<212> DNA
<213> Rattus norvegicus

<400> 22
gagtagatga tgggttca 18

EPO - Munich
83
11. Feb. 2003

Claims

1. Isolated protein comprising the same or substantially the same amino acid sequence selected from the group consisting of SEQ ID NOs: 1 to 5, or a splice variant or a salt thereof.
5
2. Protein according to claim 1, which comprises at least one fragment of said amino acid sequences.
- 10 3. Nucleic acid, which comprises at least one nucleic acid encoding a protein according to claims 1 and/or 2.
4. Nucleic acid according to claim 3, which consists of DNA or RNA.
- 15 5. Nucleic acid according to claim 4 comprising a nucleic acid selected from the group consisting of SEQ ID NOs: 6 to 10.
6. Nucleic acid according to one of claims 3 to 5 further comprising at least one promoter, enhancer, intron and/or polyA-sequence.
20
7. Nucleic acid, which is complementary to the nucleic acid according to one of claims 3 to 6.
8. Vector comprising a protein according to claims 1 or 2 and/or a nucleic acid according to one of claims 3 to 7.
25
9. Vector according to claim 8 selected from the group of vectors consisting of plasmids, phagemids, phages, cosmids, artificial mammalian chromosomes, knock-out or knock-in constructs, viruses, in particular adenovirus, vaccinia virus, baculovirus, retrovirus, adeno-associated-virus, rhinovirus, HIV, adeno-associated virus (AAV), herpes simplex virus (HSV-1), lentivirus, filovirus and engineered versions thereof, naked DNA, virosomes, liposomes, nucleic acid coated particles, in particular gold spheres.
30

10. Isolated cell comprising a protein according to claims 1 or 2, a nucleic acid according to claims 3 to 7 and/or a vector according to claims 8 or 9.
- 5 11. Cell according to claim 10, which is a stem cell, a neuronal precursor cell or a neuronal cell, in particular an axon.
12. Transgenic non-human animal generated from a cell according to claims 10 or 11.
13. Antibody directed against a protein according to claims 1 or 2.
- 10 14. Method of producing a protein according to claims 1 or 2 and/or a nucleic acid according to claims 3 to 7 comprising the steps of:
 - a) cultivating a cell according to claims 10 or 11, and
 - 15 b) isolating said protein and/or said nucleic acid.
15. Method of isolating compounds interacting with a protein according to claims 1 or 2 comprising the steps of:
 - 20 a) contacting said protein with at least one potentially interacting compound,
 - b) measuring binding of said compound to said protein.
16. Method according to claim 16, further comprising the steps of:
 - 25 a) selecting a binding compound,
 - b) modifying the binding compound, to generate a variety of modified binding compounds,
 - c) contacting said protein with each of the modified binding compounds,
 - d) measuring binding of said modified compounds to said protein, and
 - 30 e) if needed repeating steps a) to d) for one or more times.
17. Method of isolating functional interactors comprising the steps of:

- a) contacting a neuronal cell that comprises a wt nucleic acid coding for a protein selected from the group consisting of SEQ ID NOs: 1 to 5, a splice variant thereof, or a fragment thereof with a potential functional interactor,
- b) contacting the cell with a bioactive lipid phosphate, and
- c) measuring neurite movement.

18. Method according to claim 17 further comprising the steps of:

- a) contacting a neuronal cell that comprises a mutant nucleic acid coding for a mutant of the protein selected from the group consisting of SEQ ID NOs: 1 to 5, or a splice variant thereof or that contains a knock-out of the wt nucleic acid coding for one of said proteins with a potential functional interactor,
- b) contacting said cell with a bioactive lipid phosphate, and
- c) measuring neurite movement.

19. Method according to claims 17 or 18 further comprising the steps of:

- a) selecting a functional interactor,
- b) modifying the functional interactor, to generate a variety of modified functional interactors,
- c) contacting a neuronal cell as used in claim 17 a) and if needed a cell as used in claim 18 a) with each of the modified functional interactors,
- d) contacting said cell or cells with a bioactive lipid phosphate,
- e) measuring neurite movement, and
- f) if needed repeating steps a) to d) for one or more times.

20. Method according to one of claims 15 to 19, further comprising the step of admixing the interacting compound or the functional interactor with suitable auxiliary substances and/or additives.

21. Pharmaceutical composition for the treatment of neuronal injuries or diseases comprising a protein according to claims 1 or 2, a nucleic acid according to claims 3 to 7, a vector according to claims 8 to 9, a cell according to claims 10 to 11, an antibody according to claims 13, a binding compound isolated by the method of claims 15 or 16

and/or a functional interactor isolated by the method of claims 17 to 19 and if needed suitable auxiliary substances and/or additives.

- 22. Use of a pharmaceutical composition of claim 21 for the production of a medicament
5 for the treatment of neuronal diseases or injuries.
- 23. Use of a protein according to claim 1 or 2 or a nucleic acid according to claims 3 to 7
as a diagnostic marker for the diagnosis of a disease or disease state.
- 10 24. Use according to claim 23, where the disease is a neuronal disease, a tumor disease or
infertility.

Summary

EPO - Munich
63
11. Feb. 2003

- 5 Lipid phosphate phosphatase proteins, genes coding for them, vectors and cells comprising them, antibodies directed against them, methods of identifying compounds binding to them and functional interactors as well as to the use of proteins, genes, vectors, cells, interacting compounds and functional interactors for treating neuronal diseases and/or injuries.

THIS PAGE BLANK (USPTO)

Figure 1

EPO - Munich
63
11. Feb. 2003

1 10 20 30 40 50 60
MQRAGSSGGR GECDISGAGR LGLEEAARLS CAVHTSPGGG RRPQQAAGMS AKERPKGKVI
70 80 90 100 110 120
KDSVTLLPCF YFVELPILAS SVVSLYFLEL TDVFKPVHSG FSCYDRSLSM PYIEPTQEI
130 140 150 160 170 180
PFLMLLSLAF AGPAITIMVG EGILYCCLSK RRNGVGLEPN INAGGCNFNS FLRRAVRFVG
190 200 210 220 230 240
VHVFGLCSTA LITDIIQLST GYQAPYFLTV CKPNYTSNLV SCKENSYIVE DICSGSDLTV
250 260 270 280 290 300
INSGRKSFPS QHATLAFAA VYVSMYFNST LTDSSKLLKP LLVFTFIICG IICGLTRITQ
310 320 330 340 350 360
YKNHPVDVYC GFLIGGGIAL YLGLYAVGNF LPSDESMFOH RDALRS LDTL NQDENRELSA
370 380 390 400 410 420
KNGSSSDGTA HTEGILNRNH RDASSLTNLK RANADVEIT PRSPNGKENM VTFSTLPRRA
430 440 450 460 470 480
NTPSVEDPVR RNASTHASMD SARSKQULTQ WKKNKESRKL SLOVIEPEPG OSPPRSTEMR
490 500 510 520 530 540
SSSEPSRVGV NGDHHGPGNQ YLKIQPGAVP GCNNSMPGGP RVSTQSRPGS SOLVHPEET
550 560 570 580 590 600
QENISTSPKS SSARAKWLKA AEKTVAGNRS NSQPRIMQVI AMSKQOGVLO SSPKNTEGST
610 620 630 640 650 660
VSCTGSIRYK TLTDHEPSGI VRVEAHEENN RPTIOTPSTE GEGSGSNKWK APEKGSERQT
670 680 690 700 710 720
VELNDENROS LSCGSLKDSF GSGDEKESNI DSNEHHHGGT TMRVTPVEG SEIGSETIST
730 740 750 760 770 780
SSSRDSTLRR KGNITLPER SNSPENTRNI FYKGTSETRA YKD

Figure 2




Figure 3




Figure 4






Figure 5

ipsilateral hippocampus



contralateral hippocampus



cortex




Figure 6




Figure 7






Figure 8




Figure 9




Figure 10




Figure 11




Figure 12




Figure 13




Figure 14

